

ALMOST SURE MULTIFRACTAL SPECTRUM FOR THE TIP OF AN SLE CURVE

FREDRIK JOHANSSON VIKLUND AND GREGORY F. LAWLER

ABSTRACT. The tip multifractal spectrum of a two-dimensional curve is one way to describe the behavior of the uniformizing conformal map of the complement near the tip. We give the tip multifractal spectrum for a Schramm-Loewner evolution (SLE) curve, we prove that the spectrum is valid with probability one, and we give applications to the scaling of harmonic measure at the tip.

1. INTRODUCTION

The chordal Schramm-Loewner evolution (SLE_κ) is a one parameter family of probability measures on curves $\gamma : [0, \infty) \rightarrow \overline{\mathbb{H}}$, where $\mathbb{H}$ denotes the complex upper half plane. It was invented by Schramm [19] as a candidate for the scaling limit of two-dimensional lattice models from statistical physics that satisfy conformal invariance and a Markovian property in the limit. Several lattice models have since been shown to have scaling limits that can be described by SLE. Examples include loop-erased random walk and the uniform spanning tree [14], the percolation exploration-process [20], and the FK-Ising model [21]. We refer the reader to [7, 8, 22] for surveys and further references.

In this paper we will be interested in the behavior at the tip $\gamma(t)$ of the growing SLE curve. Since the curves are fractals, one cannot make sense of derivatives. Instead, the natural approach is to consider the behavior of $|g'_t(z)|$ for z near $\gamma(t)$ where g_t is a uniformizing conformal map from the complement of the curve to the upper half plane. For technical reasons, it is often easier to consider $f_t = g_t^{-1}$ near V_t , the pre-image of the tip on the real-line. Our main goal will be to derive the almost sure *tip multifractal spectrum* for SLE. For a suitable interval of α , it is defined roughly as the dimension of the subset of the curve corresponding to t for which $y|f'_t(iy + V_t)|$ decays like y^α when $y \rightarrow 0+$. We shall see that the tip multifractal spectrum is closely related to the multifractal spectrum of harmonic measure at the tip. As a function of α , this spectrum measures the size of the part of the curve that corresponds to t for which the harmonic measure of a ball of radius ϵ centered at the tip decays like ϵ^α as $\epsilon \rightarrow 0+$.

Johansson Viklund acknowledges the support of the Simons Foundation and the Knut and Alice Wallenberg Foundation.

Lawler is supported by National Science Foundation grant DMS-0907143.

The multifractal spectrum of harmonic measure has been studied extensively in the physics and mathematics literature. For example, in the case of the paths of Brownian motion, the spectrum is determined by the Brownian intersection exponents, see [11] and the references therein. In two dimensions these exponents were established by Lawler, Schramm, and Werner in [11, 12, 13]. In the case of the SLE path, Duplantier used non-rigorous “quantum gravity” methods to predict a harmonic measure spectrum for the tip, see [3, Section 7]. However, this spectrum is different to the ones we will work with as it describes the *local* dimension of harmonic measure; it corresponds in some sense to our function $\rho(\beta)$, see Section 3. (It also does not consider a “generic” tip but rather the non-equivalent behavior at the bulk point of a radial SLE path.) Using similar methods, Duplantier and Binder predicted the spectrum of harmonic measure for the *bulk* of SLE, see [4]. Roughly speaking, this spectrum is defined as the dimension of the subset of the curve away from the tip where, for a given α , harmonic measure in a shrinking ball of radius ϵ decays like ϵ^α . Beliaev and Smirnov [2] made a start to proving this result by establishing the *average integral means spectrum* for SLE. To get the almost sure multifractal spectrum from the average integral means spectrum one can formally apply the so-called multifractal formalism [16] and find the bulk spectrum by taking a Legendre transform of the average integral means spectrum. This approach is believed to be valid for SLE, although it has not been established in this case. Indeed, to the best of our knowledge, our results are the first on almost sure multifractal spectra for the family SLE_κ , $\kappa > 0$.

The starting point of our analysis is estimation of moments of the derivative of f_t using the reverse-time Loewner flow; this was started by Rohde and Schramm in [18] and extended in many places, e.g., [2, 5, 6, 9, 15]. (This is the analogue of the average integral means spectrum result for our problem.) In order to get almost sure results, one needs second moment estimates. The ideas for that appear in [9] and they were used in, e.g., [5]. These ideas are also important in understanding the so-called natural parameterization of SLE curves, see [10].

1.1. Multifractal spectra for the tip. We now proceed to discuss in more detail the multifractal spectra that we will consider. To motivate our definitions, we will start out in a slightly different setting than the one we will work with in the bulk of the paper.

Suppose that ζ is a boundary point of a simply connected domain D . We say that ζ is *accessible (in D by η)* if $\eta : [0, 1] \rightarrow \mathbb{C}$ is a simple curve with $\eta(0) = \zeta$ and $\eta(0, 1] \subset D$. If ζ is accessible by η , let h be a conformal transformation of D onto $\mathbb{C} \setminus (-\infty, 0]$ with $h(\zeta) = 0$. By $h(\zeta) = 0$, we mean $h(\eta(0+)) = 0$.

We have the following situation in mind. Let $\tilde{\gamma} : (-\infty, \infty) \rightarrow \mathbb{C}$ be a simple curve with $\tilde{\gamma}(t) \rightarrow \infty$ as $t \rightarrow \pm\infty$. For each t , we consider the “slit” plane $D_t = \mathbb{C} \setminus \tilde{\gamma}(-\infty, t]$, which is a simply connected domain whose boundary

contains $\tilde{\gamma}(t)$ and ∞ . The (*nontangential*) *tip multifractal spectrum* which we describe in this subsection is one way to describe the behavior near $\tilde{\gamma}(t)$ of the conformal map uniformizing D_t , for different values of t . Clearly, the boundary point $\tilde{\gamma}(t)$ is accessible in D_t by the curve $\eta^{(t)}(s) = \tilde{\gamma}(t+s)$.

Remark. For endpoints of slits like $\tilde{\gamma}(t)$ in D_t , there is only one possible meaning for $h(\tilde{\gamma}(t)) = 0$, but for general D a boundary point ζ might be approached from different directions that correspond to different values of $h(\zeta)$. Formally, this can be understood using *prime ends*, see, e.g., [17, Chapter 2]. In the case at hand, the curve η specifies a particular direction/prime end.

Returning to the general simply connected domain D , let $g(z) = i\sqrt{h(z)}$, where the branch of the square root is chosen so that $\sqrt{1} = 1$. Then g is a conformal transformation of D onto the upper half plane $\mathbb{H}$ with $g(0) = 0$. The map g is only unique up to composition with a Möbius transformation, that is, if $\tilde{g}$ is another such map, then

$$\tilde{g}(z) = T[g(z)],$$

where T is a Möbius transformation of $\mathbb{H}$ fixing 0. Similarly h is not unique.

Let $\eta^*(s) = g(\eta(s))$. Then $\eta^* : (0, 1] \rightarrow \mathbb{H}$ is a curve with $\eta^*(0+) = 0$. If $\eta_1^* : (0, 1] \rightarrow \mathbb{H}$ is another curve with $\eta_1^*(0+) = 0$, and $\eta_1(t) = g^{-1}(\eta_1^*(s))$, then $\eta_1(0+) = \zeta$ and ζ is accessible by η_1 . This uses the fact that the curve η exists. If ζ is an inaccessible boundary point, then the limit $\lim_{s \rightarrow 0+} \eta_1(s)$ does not exist. We say that η^* satisfies a *weak cone condition* if there is a subpower function (see Section 2.1) ψ such that for all $s > 0$,

$$|\operatorname{Re}[\eta^*(s)]| \leq \operatorname{Im}[\eta^*(s)] \psi\left(\frac{1}{\operatorname{Im}[\eta^*(s)]}\right),$$

and we say that η is *weakly nontangential* if $g \circ \eta$ satisfies a weak cone condition. It is not difficult to see that this definition is independent of the choice of g . One example of a weakly nontangential curve for D is

$$\eta(s) = g^{-1}(si), \quad 0 < s \leq 1.$$

We will use this particular curve to define the tip multifractal spectrum but the definition will be the same for any weakly nontangential curve.

Next, we let $f = g^{-1}$ so that f is a conformal transformation of $\mathbb{H}$ onto D . Since $f(is) = \eta(s)$, $s > 0$, is a simple curve, the length of $\eta(0, s]$ is given by

$$v(f; s) := \int_0^s |f'(iy)| dy. \quad (1.1)$$

A sufficient condition for the existence of a limiting $\zeta = \eta(0+)$ is that $v(f; 0+) = 0$ which is equivalent to

$$v(f; t) < \infty, \quad t > 0.$$

We can also use the plane slit by the negative real axis as uniformizing domain and write $f(w) = F(-w^2)$ where $F : \mathbb{C} \setminus (-\infty, 0] \rightarrow D$ with $F(0) =$

ζ . Then $F^{-1}(\eta(0, s]) = [0, s^2]$. In particular, the length of $F^{-1}(\eta(0, s])$ is s^2 , and the length of $f^{-1}(\eta(0, s])$ is s .

We will say that the (*nontangential*) *scaling exponent* at the boundary point ζ is θ if

$$v(f; s) \approx^* s^{2\theta}, \quad y \rightarrow 0+.$$

In particular, if $D = \mathbb{C} \setminus (-\infty, t]$, then the scaling exponent at t equals 1. (Recall that $f : \mathbb{H} \rightarrow D$ and see (2.1) for the definition of $\approx^*$.) More generally, if γ is differentiable at t , then $\theta = 1$ at t . Note that the Beurling estimates (see Lemma 2.6) imply that $\theta \leq 1$. (In fact, the same bound holds for a “limsup version” of the definition of θ .) The scaling exponent is closely related to the behavior of $|f'(iy)|$ as $y \rightarrow 0+$. Indeed, if $y|f'(iy)| \approx^* y^{1-\beta}$ for some $\beta < 1$, then (see Proposition 2.7)

$$v(f; y) \approx^* y^{1-\beta}, \quad y \rightarrow 0+$$

so that

$$\theta = \frac{1-\beta}{2}.$$

Although the definition of $v(f; y)$ depends on the choice of conformal map f , it is not hard to see that the scaling exponent θ is independent of the choice.

Returning to the curve $\tilde{\gamma}$, we can study T_θ , the set of t such that the scaling exponent of D_t at $\tilde{\gamma}(t)$ equals θ . The tip multifractal spectrum can then be defined to be either of the two functions:

$$\theta \mapsto \dim_h(T_\theta), \quad \theta \mapsto \dim_h(\tilde{\gamma}[T_\theta]),$$

where $\dim_h$ denotes Hausdorff dimension. The first function depends on the choice of parameterization of $\tilde{\gamma}$ and the second is independent of parameterization. One could also define liminf and limsup versions of this. The main goal of this paper is to compute the tip multifractal spectrum for the chordal SLE path. For technical convenience, we will use an alternative definition in terms of the behavior of $|f'(iy)|$ as $y \rightarrow 0+$ and we will use β rather than θ as our variable.

Suppose now that $\gamma = \gamma(t)$ is a curve in $\overline{\mathbb{H}}$ with $\gamma(0+) \in \mathbb{R}$. Let H_t be the unbounded connected component of $\mathbb{H} \setminus \gamma[0, t]$. One way to define the multifractal spectrum of harmonic measure at the tip is as the function

$$\alpha \mapsto \dim_h(\gamma[T_\alpha^{\text{hm}}]),$$

where

$$T_\alpha^{\text{hm}} = \{t : \text{hm}(\mathcal{B}(\gamma(t), r)) \approx^* r^\alpha, r \rightarrow 0+\}.$$

Here $\text{hm}(\cdot) = \text{hm}(\infty, \cdot, H_t)$, is the normalized harmonic measure from infinity. We will both use this definition and a slightly different (nonequivalent) definition that is more closely related to the tip multifractal spectrum that we described above. See Section 2.3 for precise definitions.

1.2. Main results. The following is part of our main result, Theorem 3.1. For the purpose of stating the theorem, let $\hat{f}_t(z) = f_t(z + V_t)$, where $f_t : \mathbb{H} \rightarrow H_t$ is the chordal SLE_κ Loewner chain and V_t is the Loewner driving function for f_t . (See Section 3 for definitions.) Further, let

$$\rho(\beta) = \frac{\kappa}{8(\beta+1)} \left[\left(\frac{\kappa+4}{\kappa} \right) (\beta+1) - 1 \right]^2,$$

and set

$$\beta_\pm = -1 + \frac{\kappa}{12 + \kappa \mp 4\sqrt{8 + \kappa}}.$$

Define

$$\Theta_\beta = \{t \in (0, 2] : y|\hat{f}'_t(iy)| \approx^* y^{1-\beta}\}.$$

(See Section 2.1 for the definition of $\approx^*$.)

Theorem (Tip multifractal spectrum). *Suppose $\kappa > 0$ and that $\beta_- \leq \beta \leq \beta_+$. For chordal SLE_κ , with probability one,*

$$\dim_h(\Theta_\beta) = \frac{2 - \rho(\beta)}{2}, \quad \dim_h[\gamma(\Theta_\beta)] = \frac{2 - \rho(\beta)}{1 - \beta},$$

See the precise statement in Theorem 3.1. (We prove more than we state here.)

Notice that we obtain Beffara's theorem on the dimension of the SLE_κ path [1] as a corollary of Theorem 3.1.

Using the tip multifractal spectrum and some additional work we can derive the almost sure spectrum for harmonic measure at the tip; see Section 2.3 for more details. Although we modify the definition of the spectrum somewhat, we prove in Theorem 3.2 the stronger almost sure version of the theorem. To state it, define

$$\alpha_\pm = \frac{1}{1 - \beta_\pm},$$

where $\beta_\pm$ are as above.

Theorem (Multifractal spectrum for harmonic measure at the tip). *Suppose $\kappa > 0$ and that $\alpha_- \leq \alpha \leq \alpha_+$. For chordal SLE_κ , with probability one,*

$$\dim_h[\gamma(\Theta_\alpha^{\text{hm}})] = \alpha \left(1 - \frac{4}{\kappa} \right) + \frac{(4 + \kappa)^2}{8\kappa} - \frac{\kappa}{8} \left(\frac{\alpha^2}{2\alpha - 1} \right). \quad (1.2)$$

In the final section of the paper we prove Theorem 3.3 which together with Theorem 3.2 and a Beurling estimate shows that the right hand side of (1.2) gives the harmonic measure spectrum for a (one-sided) version which is closer to the usual definition, but for a smaller range of α .




FIGURE 1. Multifractal spectrum of harmonic measure at the tip for SLE_κ , $\kappa = 2, 4, 6$. The maximum is the Hausdorff dimension of the curve.

1.3. Outline of the paper. Our paper is organized as follows. The next section discusses some preliminary facts. After setting some notation about asymptotics in Section 2.1, the deterministic Loewner equation is discussed in Section 2.2. Much in this subsection is standard but we have included this in order to phrase the results appropriately for our purposes. Also, we want to separate estimates that deal only with the Loewner equation itself from those that are particular to SLE. In this subsection, there are three kinds of results: those that hold for all conformal maps of $\mathbb{H}$ for which we use the letter h ; those that hold for all solutions of the chordal Loewner equation for which we use g_t and $f_t = g_t^{-1}$; and towards the end fact about solutions of the Loewner equation for driving functions that are weakly Hölder-(1/2). We also formally define the tip multifractal spectra in this section.

The main theorem is not stated in full until Section 3 where the Schramm-Loewner evolution (SLE), that is, the solution of the Loewner equation with a Brownian motion input, is discussed. From here on a value of the SLE parameter κ is fixed and a large number of κ -dependent parameters are defined. Although we do not discuss it directly, what we are doing is establishing the guess for the value of the multifractal spectrum in terms of the Legendre transform of a logarithmic moment generating function.

The basic proofs of the main theorem can be found in Section 4. This section is relatively short because it relies on estimates on the moments of the derivative some of which were established in [9, 5]; the necessary additions are proved in Section 5. Section 6 uses the *forward* Loewner flow to prove a result on the harmonic measure spectrum stated in Section 3.3. We warn the reader that some of the notation in Section 6 does not agree with that earlier and that the assumption $\kappa < 8$ is made there.

2. PRELIMINARIES

2.1. Notations. In order to avoid writing bulky expressions with ratios of logarithms, we will adopt the following notations.

We call a function $\psi : [0, \infty) \rightarrow (0, \infty)$ a *(positive) subpower function* if it is continuous, nondecreasing, and

$$\lim_{x \rightarrow \infty} x^{-u} \psi(x) = 0,$$

for all $u > 0$.

If f and g are positive functions tending to zero with y , we write

$$f(y) \approx^* g(y), \quad y \rightarrow 0+, \quad (2.1)$$

if there exists a subpower function ψ such that

$$\psi(1/y)^{-1} g(y) \leq f(y) \leq \psi(1/y) g(y), \quad y \rightarrow 0+$$

We write

$$f(y) \preccurlyeq g(y), \quad y \rightarrow 0+,$$

if

$$\limsup_{y \rightarrow 0+} \frac{\log g(y)}{\log f(y)} \leq 1,$$

and

$$f(y) \preccurlyeq_{\text{i.o.}} g(y), \quad y \rightarrow 0+,$$

if

$$\liminf_{y \rightarrow 0+} \frac{\log g(y)}{\log f(y)} \leq 1.$$

Here i.o. stands for “infinitely often”. Clearly $f(y) \preccurlyeq g(y)$ implies $f(y) \preccurlyeq_{\text{i.o.}} g(y)$ but the converse is not true. Similarly we write $f(y) \succcurlyeq g(y)$ and $f(y) \succcurlyeq_{\text{i.o.}} g(y)$ for

$$\liminf_{y \rightarrow 0+} \frac{\log g(y)}{\log f(y)} \geq 1 \quad \text{and} \quad \limsup_{y \rightarrow 0+} \frac{\log g(y)}{\log f(y)} \geq 1,$$

respectively. We write $f(y) \approx g(y)$ if $f(y) \preccurlyeq g(y)$ and $f(y) \succcurlyeq g(y)$, that is, if

$$\lim_{y \rightarrow 0+} \frac{\log g(y)}{\log f(y)} = 1.$$

Note that if $\beta > 0$, then

$$f(y) \approx y^\beta \iff f(y) \approx^* y^\beta.$$

We will also use the notations for asymptotics for functions $f(n), g(n)$ as $n \rightarrow \infty$ along the positive integers.

2.2. Chordal Loewner equation. In this section, we review some facts about conformal mappings and the chordal Loewner equation. See [7, Chapters 3,4] for proofs of theorems stated without proof here.

Suppose $\tilde{\gamma} : (-\infty, \infty) \rightarrow \mathbb{C}$ is a curve as in the introduction. The chordal Loewner equation is an equation that describes the evolution of $\tilde{\gamma}(0, \infty)$ given $\tilde{\gamma}(-\infty, 0]$. Let $\tilde{g}$ be a conformal transformation of $\mathbb{C} \setminus \tilde{\gamma}(-\infty, 0]$ onto the upper half plane $\mathbb{H}$ with $\tilde{g}(\gamma(0)) = 0$, $\tilde{g}(\infty) = \infty$. In order to describe $\tilde{\gamma}(t), t > 0$, it suffices to describe

$$\gamma(t) := \tilde{g}(\tilde{\gamma}(t)), \quad 0 \leq t < \infty,$$

and this is what the Loewner equation in $\mathbb{H}$ does. For the remainder of the paper, we will consider a curve γ in $\mathbb{H}$ as above. The Riemann mapping theorem implies that there is a unique conformal transformation g_t of $\mathbb{H} \setminus \gamma(0, t]$ onto $\mathbb{H}$ with $g_t(z) = z + o(1)$ as $z \rightarrow \infty$. We can expand g_t at infinity,

$$g_t(z) = z + \frac{a(t)}{z} + O(|z|^{-2}),$$

where $a(t)$ by definition is the half-plane capacity of $\gamma(0, t]$. It is continuous and strictly increasing. We make the (slightly) stronger assumption that $a(t) \rightarrow \infty$ as $t \rightarrow \infty$. Then the chordal Loewner integral equation states that

$$g_t(z) = z + \int_0^t \frac{da(s)}{g_s(z) - V_s}, \quad t \leq T_z,$$

where $V_s = g_s(\gamma(s))$ and $T_z = \inf\{t : \text{Im}[g_t(z)] = 0\} = \inf\{t : g_t(z) - V_t = 0\}$. It can be shown that $s \mapsto V_s$ is a continuous function. It is convenient to choose a parameterization of γ such that $a(t) = at$ for some $a > 0$ in which case we get the Loewner differential equation

$$\partial_t g_t(z) = \frac{a}{g_t(z) - V_t}, \quad g_0(z) = z. \quad (2.2)$$

Let

$$f_t(z) = g_t^{-1}(z), \quad \hat{f}_t(z) = f_t(z + V_t) = g_t^{-1}(z + V_t).$$

By differentiating both sides of $f_t(g_t(z)) = z$ with respect to t we see that $(f'_t(z) := \partial_z f_t(z))$

$$\partial_t f_t(z) = -f'_t(z) \frac{a}{z - V_t}, \quad (2.3)$$

and since $g_t(\gamma(t)) = V_t$, we get

$$\gamma(t) = f_t(V_t) = \lim_{y \rightarrow 0+} f_t(V_t + iy) = \lim_{y \rightarrow 0+} \hat{f}_t(iy). \quad (2.4)$$

We let

$$v_t(y) = v(\hat{f}_t; y) = \int_0^y |\hat{f}'_t(iu)| du.$$

As mentioned before, if $v_t(y) < \infty$ for some $y > 0$, then $v_t(0+) = 0$ and the limit in (2.4) exists. More work is needed to determine whether or not γ is

a continuous function of t . Note that if g_t satisfies (2.2) and $g_t^* = g_{t/a}$, then

$$\partial_t g_t^*(z) = \frac{1}{g_t^*(z) - V_t^*}, \quad g_0^*(z) = z,$$

where $V_t^* = V_{t/a}$.

Conversely, we can start with a continuous function $t \mapsto V_t$ and $a > 0$ and define a Loewner chain $(g_t, t \geq 0)$ by (2.2). We define $\gamma(t)$ by (2.4) provided that the limit exists. We say that the family of conformal maps g_t *generates a curve* if γ exists and is a continuous function of t . We do *not* assume that the curve is simple. If H_t denotes the unbounded component of $\mathbb{H} \setminus \gamma(0, t]$, then g_t is the unique conformal transformation of H_t onto $\mathbb{H}$ satisfying

$$g_t(z) = z + \frac{at}{z} + O(|z|^{-2}), \quad z \rightarrow \infty.$$

Lemma 2.1. *For every t and every $y > 0$ with $v_t(y) < \infty$,*

$$\frac{y |\hat{f}'_t(iy)|}{4} \leq |\gamma(t) - \hat{f}_t(iy)| \leq v_t(y). \quad (2.5)$$

Proof. The second estimate is immediate from the definition of $v_t(y)$ and the first inequality follows from the Koebe (1/4)-theorem applied to $\hat{f}_t$ on the open disk of radius y about iy . $\square$

Lemma 2.2. *If f_t satisfies (2.3) and $z = x + iy \in \mathbb{H}$, then for $s \geq 0$*

$$e^{-5as/y^2} |f'_t(z)| \leq |f'_{t+s}(z)| \leq e^{5as/y^2} |f'_t(z)|. \quad (2.6)$$

In particular, if $s \leq y^2$,

$$e^{-5a} |f'_t(z)| \leq |f'_{t+s}(z)| \leq e^{5a} |f'_t(z)|.$$

Proof. Without loss of generality we may assume that $a = 1$. Differentiating (2.3) with respect to z yields

$$\partial_t f'_t(z) = -f''_t(z) \frac{1}{z - V_t} + f'_t(z) \frac{1}{(z - V_t)^2}.$$

Note that $|z - V_t| \geq y$. Applying Bieberbach's theorem (the $n = 2$ case of the Bieberbach conjecture) to the disk of radius y about z , we can see that

$$|f''_t(z)| \leq 4y^{-1} |f'_t(z)|.$$

and hence

$$|\partial_t f'_t(z)| \leq 5y^{-2} |f'_t(z)|,$$

which implies (2.6). $\square$

The Koebe distortion and growth theorems are traditionally stated in terms of univalent functions defined on the unit disk, see, e.g., [17, Chapter 2]. We will use these theorems for univalent functions on $\mathbb{H}$, and the next proposition gives the appropriate results.

Proposition 2.3. *Suppose $h : \mathbb{H} \rightarrow \mathbb{C}$ is a conformal transformation, $x \in \mathbb{R}$, $y > 0$, and $r \geq 1$. Then*

$$(x^2 + 4)^{-2} |h'(iy)| \leq |h'(y(x+i))| \leq (x^2 + 4)^2 |h'(iy)|, \quad (2.7)$$

$$|h(y(x+i)) - h(iy)| \leq \frac{(x^2 + 4)^{3/2} |x|}{2} y |h'(iy)|, \quad (2.8)$$

$$r^{-3} |h'(iy)| \leq |h'(iyr)| \leq r |h'(iy)|, \quad (2.9)$$

$$|h(iyr) - h(iy)| \leq \frac{r^2 - 1}{2} y |h'(iy)|. \quad (2.10)$$

Proof. By scaling we may assume that $y = 1$. Let

$$G(z) = \frac{z-i}{z+i}, \quad G'(z) = \frac{2i}{(z+i)^2},$$

which is a conformal transformation of $\mathbb{H}$ onto the unit disk $\mathbb{D}$ with $G(i) = 0$, $|G'(i)| = 1/2$. We can write

$$h(z) = f(G(z)), \quad h'(z) = f'(G(z)) G'(z),$$

where f is a univalent function on $\mathbb{D}$. The distortion theorem tells us that

$$|f'(w)| \leq \frac{1+|w|}{(1-|w|)^3} |f'(0)|, \quad |w| < 1,$$

and the growth theorem states that

$$|f(w) - f(0)| \leq \frac{|w|}{(1-|w|)^2} |f'(0)|, \quad |w| < 1.$$

Since $|G'(i)| = 1/2$, we get

$$|h'(z)| \leq \frac{2|G'(z)| [1+|G(z)|]}{(1-|G(z)|)^3} |h'(i)|, \quad (2.11)$$

and

$$|h(z) - h(i)| \leq \frac{2|G(z)|}{(1-|G(z)|)^2} |h'(i)|. \quad (2.12)$$

Since

$$|G(x+i)| = \frac{|x|}{\sqrt{x^2+4}}, \quad |G'(x+i)| = \frac{2}{x^2+4},$$

we plug into (2.11) and see that

$$|h'(x+1)| \leq \frac{1}{16} [\sqrt{x^2+4} + |x|]^4 |h'(i)| \leq (x^2+4)^2 |h'(i)|.$$

This gives the second inequality in (2.7) and the first follows easily by real translation. Plugging into (2.12) gives

$$\begin{aligned} |h(x+i) - h(i)| &\leq \frac{2|x| \sqrt{x^2+4} (\sqrt{x^2+4} + x)^2}{16} |h'(i)| \\ &\leq \frac{(x^2+4)^{3/2} |x|}{2} |h'(i)|. \end{aligned}$$

Since $r \geq 1$,

$$|G(ir)| = \frac{r-1}{1+r} = |G(i/r)|, \quad |G'(ir)| = \frac{2}{(1+r)^2}, \quad |G'(i/r)| = \frac{2r^2}{(r+1)^2}.$$

Plugging into (2.11) and (2.12) gives (2.9) and (2.10). $\square$

Corollary 2.4. *If $h : \mathbb{H} \rightarrow \mathbb{C}$ is a conformal transformation, then for every $y > 0$,*

$$v(h; y) \geq \frac{y |h'(iy)|}{2}, \quad (2.13)$$

$$\frac{2}{3} v(h; 2^{-n}) \leq \sum_{j=n}^{\infty} 2^{-j} |h'(i2^{-j})| \leq \frac{8}{3} v(h; 2^{-n}).$$

Proof. We write

$$v(h; 2^{-n}) = \sum_{j=n}^{\infty} \int_{2^{-j-1}}^{2^{-j}} |h'(iy)| dy.$$

Using (2.9) (which holds for $r > 1$)

$$\begin{aligned} v(h; y) &= \int_0^y |h'(is)| ds = y \int_0^1 |h'(iry)| dr \geq y |h'(iy)| \int_0^1 r dr = \frac{y |h'(iy)|}{2}, \\ \int_{r/2}^r |h'(iy)| dy &\leq r |h'_t(ir)| \int_{1/2}^1 s^{-3} ds = \frac{3r}{2} |h'(ir)|, \\ \int_{r/2}^r |h'(iy)| dy &\geq r |h'(ir)| \int_{1/2}^1 s ds = \frac{3r}{8} |h'(ir)|. \end{aligned}$$

$\square$

We define the following measure of the modulus of continuity of U_t :

$$\Delta(t, s) = \sup_{0 \leq r \leq s^2} \sqrt{s^{-2}(U_{t+r} - U_t)^2 + 4}.$$

Note that $\Delta(t, s) \geq 2$ and it is of order one if

$$\sup_{0 \leq r \leq s^2} |U_{t+r} - U_t| \approx s.$$

The definition of $\Delta(t, s)$ with the 4 has been chosen to make the statement of the next proposition cleaner.

Proposition 2.5. *If $t \geq 0$ and $0 \leq r \leq s^2$ with $v_t(s) + v_{t+r}(s) < \infty$, then*

$$|\gamma(t+r) - \gamma(t)| \leq v_t(s) + v_{t+r}(s) + e^{5a} |\hat{f}'_t(si)| \Delta(t, s)^4 s. \quad (2.14)$$

Proof. By the triangle inequality and (2.5),

$$\begin{aligned} |\gamma(t+r) - \gamma(t)| &\leq |\hat{f}_t(si) - \gamma(t)| + |\hat{f}_{t+r}(si) - \gamma(t+r)| + |\hat{f}_t(si) - \hat{f}_{t+r}(si)| \\ &\leq v_t(s) + v_{t+r}(s) + |f_{t+r}(V_{t+r} + si) - f_t(V_t + si)|. \end{aligned}$$

Also,

$$|f_{t+r}(V_{t+r} + si) - f_t(V_t + si)| \leq |f_{t+r}(V_{t+r} + si) - f_{t+r}(V_t + si)| + |f_{t+r}(V_t + si) - f_t(V_t + si)|.$$

Using (2.8) and (2.6), we see that

$$\begin{aligned} |f_{t+r}(V_{t+r} + si) - f_{t+r}(V_t + si)| &\leq \frac{1}{2} |f'_{t+r}(V_t + si)| \Delta(t, s)^4 s \\ &\leq \frac{1}{2} e^{5a} |f'_t(V_t + si)| \Delta(t, s)^4 s \\ &= \frac{1}{2} e^{5a} |\hat{f}'_t(si)| \Delta(t, s)^4 s \end{aligned}$$

Also (2.6) and (2.3) imply that

$$|\partial_r f_{t+r}(V_t + is)| \leq \frac{a}{s} e^{5ar/s^2} |\hat{f}'_t(is)|, \quad (2.15)$$

and hence

$$\begin{aligned} |f_{t+r}(V_t + is) - f_t(V_t + is)| &\leq |f'_t(V_t + is)| \int_0^{s^2} \frac{a}{s} e^{5au/s^2} du \\ &= \frac{s}{5} e^{5a} |\hat{f}'_t(is)| \end{aligned} \quad (2.16)$$

$$< \frac{s}{2} e^{5a} \Delta(t, s)^4 |\hat{f}'_t(is)|. \quad (2.17)$$

□

Lemma 2.6. *There exist $c > 0$ such that for $t \geq 0$ and $0 < y \leq 1$,*

$$c \frac{y}{\sqrt{2at+1}} \leq |\hat{f}'_t(iy)| \leq \frac{\sqrt{2at+1}}{y}.$$

Proof. We may assume $a = 1$ for otherwise we consider $g_t^* = g_{t/a}$. Let $w = \hat{f}_t(iy)$, that is, $g_t(w) = V_t + iy$, and let $Y_t = \text{Im}[g_t(w)]$. The Loewner equation implies that $\partial_t[Y_t^2] \geq -2$ and hence $\text{Im}[w] \leq \sqrt{2t+1}$. Similarly, $\text{Im}[\gamma(s)] \leq \sqrt{2t} \leq \sqrt{2t+1}$ for $0 \leq s \leq t$. The Loewner equation also implies that $\partial_t[Y_t/|g'_t(w)|] \leq 0$, which implies

$$y |\hat{f}'_t(iy)| = \frac{Y_t}{|g'_t(w)|} \leq \frac{Y_0}{|g'_0(w)|} = \text{Im}[w] \leq \sqrt{2t+1}.$$

This gives the second inequality.

For the first inequality, let $d = \text{dist}(w, \gamma[0, t] \cup \mathbb{R})$. The Beurling estimate [7, Theorem 3.76] implies that there is a $c_* < \infty$ such that the probability that a Brownian motion starting at w goes distance $\sqrt{2t+1}$ without hitting $\gamma[0, t] \cup \mathbb{R}$ is bounded above by

$$c_* \left(\frac{d}{\sqrt{2t+1}} \right)^{1/2}.$$

By the gambler's ruin estimate, the probability that a Brownian motion in $\mathbb{H}$ starting at iy reaches $I_t := \{\tilde{w} : \text{Im}[\tilde{w}] = 2\sqrt{2t+1}\}$ before hitting the real

line equals $y [2\sqrt{2t+1}]^{-1}$. Since the imaginary part decreases in the forward Loewner flow, it follows from conformal invariance that the probability that a Brownian motion starting at w reaches I_t before hitting $\gamma[0, t] \cup \mathbb{R}$ is at least $y [2\sqrt{2t+1}]^{-1}$. Therefore

$$c_* \left(\frac{d}{\sqrt{2t+1}} \right)^{1/2} \geq \frac{y}{2\sqrt{2t+1}}.$$

The Koebe (1/4)-theorem implies that $d \leq 4y |\hat{f}'_t(iy)|$, and plugging in we get

$$|\hat{f}'_t(iy)| \geq \frac{y}{16 c_*^2 \sqrt{2t+1}}.$$

□

Proposition 2.7. *Suppose $h : \mathbb{H} \rightarrow \mathbb{C}$ is a conformal transformation and $v(h; y)$ is as defined in (1.1). Then for every $\beta < 1$, as $y \rightarrow 0+$*

$$\begin{aligned} y|h'(iy)| &\preceq y^{1-\beta} \text{ if and only if } v(h; y) \preceq y^{1-\beta}, \\ y|h'(iy)| &\succ_{\text{i.o.}} y^{1-\beta} \text{ if and only if } v(h; y) \succ_{\text{i.o.}} y^{1-\beta}. \\ y|h'(iy)| &\approx^* y^{1-\beta} \text{ if and only if } v(h; y) \approx^* y^{1-\beta}. \end{aligned} \quad (2.18)$$

Proof. Using Corollary 2.4, all of the assertions follow easily except that $v(h; y) \approx^* y^{1-\beta}$ implies $y|h'(iy)| \approx^* y^{1-\beta}$ which we will show here. Assume $v(h; y) \approx^* y^{1-\beta}$. By (2.13), we know that $y|h'(iy)| \preceq y^{1-\beta}$. For $0 < \epsilon < 1-\beta$, let

$$u = u_\epsilon = \frac{3\epsilon}{1-\beta-\epsilon},$$

and note that (2.9) implies for y sufficiently small

$$\begin{aligned} y^{1-\beta+\epsilon} \leq v(h; y) &= v(h; y^{1+u}) + \int_{y^{1+u}}^y |h'(is)| ds \\ &\leq y^{1-\beta+2\epsilon} + \int_{y^{1+u}}^y |h'(is)| ds \\ &\leq y^{1-\beta+2\epsilon} + y^{1-3u} |h'(iy)|. \end{aligned}$$

Hence for all y sufficiently small,

$$y|h'(iy)| \geq \frac{1}{2} y^{1-\beta} y^{3u+\epsilon}.$$

Since $u_\epsilon \rightarrow 0$ as $\epsilon \rightarrow 0+$, this gives $y|h'(iy)| \succ y^{1-\beta}$. □

Definition. For every $-1 \leq \beta \leq 1$, let

$$\begin{aligned} \overline{\Theta}_\beta &= \{t \in (0, 2] : y|\hat{f}'_t(iy)| \succ_{\text{i.o.}} y^{1-\beta}\}, \\ \Theta_\beta &= \{t \in (0, 2] : y|\hat{f}'_t(iy)| \approx^* y^{1-\beta}\}, \\ \tilde{\Theta}_\beta &= \{t \in (0, 2] : y|\hat{f}'_t(iy)| \preceq y^{1-\beta}\}, \\ \underline{\Theta}_\beta &= \{t \in (0, 2] : y|\hat{f}'_t(iy)| \preceq_{\text{i.o.}} y^{1-\beta}\}, \\ \underline{\Theta}_\beta^* &= \{t \in (0, 2] : v_t(y) \preceq_{\text{i.o.}} y^{1-\beta}\}, \end{aligned}$$

where in each case the asymptotics are as $y \rightarrow 0+$.

If $\beta \neq 1$, we can write these sets as the set of $t \in (0, 2]$ such that

$$\limsup_{y \rightarrow 0+} \frac{\log |\hat{f}'_t(iy)|}{\log(1/y)} \geq \beta,$$

$$\lim_{y \rightarrow 0+} \frac{\log |\hat{f}'_t(iy)|}{\log(1/y)} = \beta,$$

$$\limsup_{y \rightarrow 0+} \frac{\log |\hat{f}'_t(iy)|}{\log(1/y)} \leq \beta,$$

$$\liminf_{y \rightarrow 0+} \frac{\log |\hat{f}'_t(iy)|}{\log(1/y)} \leq \beta,$$

$$\liminf_{y \rightarrow 0+} \frac{\log v_t(y)}{\log(1/y)} \leq \beta - 1,$$

respectively.

Using Lemma 2.6, we can see that for every $\beta > 1$,

$$\overline{\Theta}_\beta = \Theta_\beta = \Theta_{-\beta} = \tilde{\Theta}_{-\beta} = \underline{\Theta}_{-\beta} = \emptyset.$$

Note that (2.13) implies that $\underline{\Theta}_\beta^* \subset \underline{\Theta}_\beta$. Using Proposition 2.7 we can see that we can also write

$$\Theta_\beta = \{t \in (0, 2] : v_t(y) \approx^* y^{1-\beta}, \ y \rightarrow 0+\},$$

and similarly for $\overline{\Theta}_\beta, \tilde{\Theta}_\beta$. Also, $\overline{\Theta}_\beta \cup \tilde{\Theta}_\beta = (0, 2]$ and

$$\Theta_\beta \subset \overline{\Theta}_\beta \cap \tilde{\Theta}_\beta \cap \underline{\Theta}_\beta^*.$$

Definition. The driving function V_t is *weakly Hölder-(1/2)* on $[0, 2]$ if for each $\alpha < 1/2$, V_t is Hölder continuous of order α on $[0, 2]$.

Two equivalent definitions are

- If

$$\delta(s) = \sup\{|V_{t+s} - V_t| : 0 \leq t \leq t+s \leq 2\},$$

then

$$\psi(x) = \sup_{s \geq 1/x} s^{-1/2} \delta(s)$$

is a subpower function.

- There is a subpower function ψ such that for all $0 \leq t \leq 2, 0 \leq s \leq 1$,

$$\Delta(t, s) \leq \psi(1/s).$$

The next proposition shows that for weakly Hölder-(1/2) functions V_t , it suffices to consider dyadic times in the definition of Θ_β , etc.

Proposition 2.8. *Suppose V_t is weakly Hölder-(1/2) on $[0, 2]$. For each $t \in [0, 2]$ define $t_n = t_n(t) = (j-1)/2^{2n}$ if*

$$\frac{j-1}{2^{2n}} \leq t < \frac{j}{2^{2n}}.$$

Then for $-1 \leq \beta \leq 1$,

•

$$\Theta_\beta = \left\{ t \in (0, 2] : 2^{-n} |\hat{f}'_{t_n}(i2^{-n})| \approx^* 2^{-n(1-\beta)} \right\},$$

$$\bar{\Theta}_\beta = \left\{ t \in (0, 2] : 2^{-n} |\hat{f}'_{t_n}(i2^{-n})| \succ_{\text{i.o.}} 2^{-n(1-\beta)} \right\},$$

$$\underline{\Theta}_\beta = \left\{ t \in (0, 2] : 2^{-n} |\hat{f}'_{t_n}(i2^{-n})| \preccurlyeq_{\text{i.o.}} 2^{-n(1-\beta)} \right\},$$

where the asymptotics are as $n \rightarrow \infty$ along the integers.

• *If $t \in \bar{\Theta}_\beta$,*

$$v_t(y) \succ_{\text{i.o.}} y^{1-\beta}, \quad |\gamma(t) - \hat{f}_t(iy)| \succ_{\text{i.o.}} y^{1-\beta}, \quad y \rightarrow 0+.$$

• *If $t \in \tilde{\Theta}_\beta$,*

$$v_t(y) \preccurlyeq y^{1-\beta}, \quad |\gamma(t) - \hat{f}_t(iy)| \preccurlyeq y^{1-\beta}, \quad y \rightarrow 0+. \quad (2.19)$$

• *If $t \in \Theta_\beta$,*

$$v_t(y) \approx^* y^{1-\beta}, \quad |\gamma(t) - \hat{f}_t(iy)| \approx^* y^{1-\beta}, \quad y \rightarrow 0+.$$

Proof. Note that

$$\begin{aligned} |\hat{f}'_t(i2^{-n})| &= |f'_t(V_t + i2^{-n})| \leq \Delta(t, 2^{-n})^4 |f'_t(V_t + i2^{-n})| \\ &\leq e^{5a} \Delta(t, 2^{-n})^4 |\hat{f}'_{t_n}(i2^{-n})|, \end{aligned}$$

and similarly

$$|\hat{f}'_t(i2^{-n})| \geq e^{-5a} \Delta(t, 2^{-n})^{-4} |\hat{f}'_{t_n}(i2^{-n})|.$$

Hence if V_t is weakly Hölder-(1/2), there is a subpower function ψ such that for all t, n ,

$$\psi(2^n)^{-1} |\hat{f}'_{t_n}(i2^{-n})| \leq |\hat{f}'_t(i2^{-n})| \leq \psi(2^n) |\hat{f}'_{t_n}(i2^{-n})|.$$

This implies the first assertion and the remaining follow from (2.5). $\square$

2.3. Harmonic measure at the tip. We will now discuss harmonic measure giving two nonequivalent definitions, one that is standard and one which is more directly related to the multifractal spectrum we have discussed.

In this subsection γ denotes a curve in $\mathbb{H}$ with one endpoint on the real line. We assume that the curve comes from a Loewner chain driven by a continuous function V_t , so it may have double points but it does not cross itself. Let H_t be the unbounded connected component of $\mathbb{H} \setminus \gamma[0, t]$. As above, we write $g_t : H_t \rightarrow \mathbb{H}$ for the normalized conformal mapping so that $\lim_{y \rightarrow 0+} \hat{f}_t(iy) = \gamma(t)$, where $f_t = g_t^{-1}$ and $\hat{f}_t(z) = f_t(z + V_t)$. If the curve




FIGURE 2. Sketch showing the two different subsets of the boundary that we use to define $\tilde{\Theta}_\alpha^{\text{hm}}$ (left) and $\Theta_\alpha^{\text{hm}}$ (right).

has double points, we are interpreting $\gamma(t)$ in terms of prime ends, and we then tacitly understand $\gamma(t)$ as the prime end corresponding to V_t .

If $z \in H_t$, then $\text{hm}_{t,z}$ will denote the usual harmonic measure of $\mathbb{R} \cup \gamma(0, t]$ starting at z , that is to say the hitting measure of Brownian motion starting at z stopped when it reaches ∂H_t . We let

$$\text{hm}_t(U) = \lim_{y \rightarrow \infty} y \text{hm}_{t,iy}(U),$$

which is the normalized harmonic measure from the boundary point at infinity. Note that for each $z \in H_t$, hm_t and $\text{hm}_{t,z}$ are mutually absolutely continuous. Also, conformal invariance, the normalization at infinity, and the well-known Poisson kernel in $\mathbb{H}$ show that for bounded U ,

$$\text{hm}_t(U) = \frac{1}{\pi} \text{length}[g_t(U)].$$

Let

$$\tilde{\mu}(t, \epsilon) = \text{hm}_t[\overline{\mathcal{B}}(\gamma(t), \epsilon)],$$

where $\mathcal{B}(z, \epsilon)$ denotes the open disk of radius ϵ about z with closure $\overline{\mathcal{B}}(z, \epsilon)$. For $\alpha > 0$, define

$$\tilde{\Theta}_\alpha^{\text{hm}} = \{t \in (0, 2] : \tilde{\mu}(t, \epsilon) \approx^* \epsilon^\alpha, \epsilon \rightarrow 0+\}.$$

We define the multifractal spectrum of harmonic measure at the tip by

$$\alpha \longmapsto \dim_h \left[\gamma(\tilde{\Theta}_\alpha^{\text{hm}}) \right].$$

This multifractal spectrum can be hard to compute. One of the difficulties is that $\mathcal{B}(\gamma(t), \epsilon) \cap H_t$ can contain many connected components whose images under g_t are far apart. We will give a different definition that is more directly related to the tip multifractal spectrum in this paper.

Fix $t > 0, \epsilon > 0$ and let $\mathcal{B} = \mathcal{B}(\gamma(t), \epsilon)$. Let $O = O_{t,\epsilon}$ denote the connected component of $\mathcal{B} \cap H_t$ that contains $\gamma(t)$ (considered as a prime end) on its

boundary. There is a connected component (open arc), $\sigma = \sigma_{t,\epsilon}$ of $\partial\mathcal{B} \cap H_t$ that is in ∂O and such that every curve from $\gamma(t)$ (again viewed as a prime end) to infinity in H_t passes through σ . We require also that σ separates all other such open arcs from infinity in H_t ; σ is then unique.

Let $x_- = x_{-,t,\epsilon} < V_t < x_+ = x_{+,t,\epsilon}$ denote the images of the endpoints of σ under g_t (which always exist, see, e.g., [17]). Let $E = E_{t,\epsilon}$ denote the preimage of $[x_-, x_+]$ under g_t , and let

$$\mu(t, \epsilon) = \text{hm}_t[E] = \frac{x_+ - x_-}{\pi}.$$

It is not necessarily true that $E \subset \overline{\mathcal{B}}(\gamma(t), \epsilon)$. However an estimate using the Beurling projection theorem shows that there is a $c < \infty$ such that

$$\mu(t, \epsilon/2) \leq c \tilde{\mu}(t, \epsilon). \quad (2.20)$$

We define

$$\Theta_\alpha^{\text{hm}} = \{t \in (0, 2] : \mu(t, \epsilon) \approx^* \epsilon^\alpha, \epsilon \rightarrow 0+\}.$$

The next lemma makes the connection with the tip multifractal spectrum.

Lemma 2.9. *If $1/2 \leq \alpha < \infty$, then*

$$\Theta_\alpha^{\text{hm}} = \Theta_{1-1/\alpha}.$$

Proof. We will prove that there exist $0 < c_1, c_2 < \infty$ such that for all $t \geq 0$ and all $\epsilon > 0$ sufficiently small,

$$\mu(t, 2v_t(\epsilon)) \geq c_1\epsilon, \quad (2.21)$$

$$\mu(t, \epsilon) |\hat{f}_t'(i\mu(t, \epsilon))| \leq c_2\epsilon. \quad (2.22)$$

The lemma follows immediately from these estimates combined with Proposition 2.6.

Let η_ϵ denote the line segment $(0, i\epsilon]$. The harmonic measure from infinity of η_ϵ in $\mathbb{H} \setminus \eta_\epsilon$ equals $c_1\epsilon$ for a specific constant c_1 , and hence by conformal invariance the harmonic measure from infinity of $\eta_\epsilon^* := \hat{f}_t \circ \eta_\epsilon$ in $H_t \setminus \eta_\epsilon^*$ is also $c_1\epsilon$. Since η_ϵ^* is a curve of length $v_t(\epsilon)$ and one of its endpoints is $\gamma(t)$, the interior of η_ϵ^* is contained in $O_{t, v_t(\epsilon)}$. From this and a Beurling estimate as in (2.20), we get (2.21).

It remains to prove (2.22). To this end, let $\sigma_\epsilon = \sigma_{\epsilon, t}$ be the open arc whose endpoints are mapped to $x_{-, \epsilon} < x_{+, \epsilon}$ as above. Let $\ell_\epsilon = x_{+, \epsilon} - x_{-, \epsilon}$ and note that $\mu(t, \epsilon) = \ell_\epsilon / \pi$. As $\epsilon \rightarrow 0+$, $\ell_\epsilon \rightarrow 0$ (using, e.g., the Beurling estimate), and hence for y sufficiently small we can choose ϵ such that $\ell_\epsilon \leq y$ and $\ell_{2\epsilon} \geq y$. Hence it suffices to show that $2\epsilon \geq c y |\hat{f}_t'(iy)|$. Since $\ell_{2\epsilon} \geq y$, we can see that there exists c_2 such that the probability that a Brownian motion starting at iy hits $g_t \circ \sigma_{2\epsilon}$ before leaving $\mathbb{H}$ is at least c_2 . By conformal invariance, this is also true for a Brownian motion starting at $\hat{f}_t(iy)$ hitting $\sigma_{2\epsilon}$ before leaving H_t . The distortion theorem and the Koebe-(1/4) theorem show that $\text{dist}[\hat{f}_t(iy), \partial H_t] \asymp y |\hat{f}_t'(iy)|$. The needed estimate then comes

from the Beurling estimate which implies in any simply connected domain D , if $V \subset \partial D$,

$$\text{hm}_D(z, V) \leq c \sqrt{\frac{\text{diam}(V)}{\text{dist}(z, \partial D)}}.$$

□

3. TIP SPECTRUM FOR SLE

Let $\kappa > 0$ and $a = 2/\kappa$. Then the chordal Schramm-Loewner evolution with parameter κ (SLE_κ) is the solution to (2.2) with $a = 2/\kappa$ where V_t is a standard Brownian motion. It is well known that with probability one, V_t is weakly Hölder-(1/2). Let

$$d = \min \left\{ 1 + \frac{\kappa}{8}, 2 \right\}.$$

It was proved by Beffara [1] that d is the Hausdorff dimension of the path $\gamma[0, 2]$. This will follow as a particular case of our main theorem, so we will not need to assume this result. However, it is convenient to use this notation.

3.1. Main theorem. Before stating the main theorem, we will define some special values of the parameter β . See Section 3.4 for more details. Let

$$\rho(\beta) = \frac{\kappa}{8(\beta+1)} \left[\left(\frac{\kappa+4}{\kappa} \right) (\beta+1) - 1 \right]^2, \quad (3.1)$$

$$\hat{d}_\beta = \frac{2 - \rho(\beta)}{2},$$

$$d_\beta = \frac{2\hat{d}_\beta}{1-\beta} = \frac{2 - \rho(\beta)}{1-\beta}.$$

The maximum value of $\hat{d}_\beta$ equals 1 and is obtained at

$$\beta_\# := \frac{\kappa}{\kappa+4} - 1.$$

The maximum value of d_β equals d and is obtained at

$$\beta_* := \frac{\kappa}{\max\{4, \kappa-4\}} - 1.$$

We define $\beta_- \leq \beta_\# \leq \beta_* \leq \beta_+$ by $\rho(\beta_-) = \rho(\beta_+) = 2$. A straightforward computation gives

$$\beta_+ = -1 + \frac{\kappa}{12 + \kappa - 4\sqrt{8 + \kappa}}, \quad (3.2)$$

$$\beta_- = -1 + \frac{\kappa}{12 + \kappa + 4\sqrt{8 + \kappa}} < 0. \quad (3.3)$$

Also $-1 < \beta_- < \beta_+ \leq 1$ with equality only for $\kappa = 8$.

Remark. The function $\beta_+(\kappa)$ determines the optimal Hölder exponent for the SLE_κ path in the capacity parameterization: With probability one, the chordal SLE_κ path away from the base is Hölder- α for $\alpha < (1 - \beta_+)/2$ and not Hölder- α for $\alpha > (1 - \beta_+)/2$. See Theorem 1.1 of [5] for a precise statement.

Theorem 3.1. *For chordal SLE_κ , if $-1 \leq \beta \leq 1$, the following holds with probability one.*

- If $\beta_- \leq \beta \leq \beta_+$,

$$\dim_h(\Theta_\beta) = \hat{d}_\beta, \quad \dim_h[\gamma(\Theta_\beta)] = d_\beta. \quad (3.4)$$

- If $\beta_\# \leq \beta \leq \beta_+$,

$$\dim_h(\overline{\Theta}_\beta) = \hat{d}_\beta. \quad (3.5)$$

- If $\beta_* \leq \beta \leq \beta_+$,

$$\dim_h[\gamma(\overline{\Theta}_\beta)] = d_\beta. \quad (3.6)$$

- If $\beta_- \leq \beta \leq \beta_\#$,

$$\dim_h(\underline{\Theta}_\beta) = \hat{d}_\beta. \quad (3.7)$$

- If $\beta_- \leq \beta \leq \beta_*$,

$$\dim_h[\gamma(\underline{\Theta}_\beta^*)] = d_\beta. \quad (3.8)$$

- If $\beta > \beta_+$, $\overline{\Theta}_\beta = \emptyset$.
- If $\beta < \beta_-$, $\underline{\Theta}_\beta = \emptyset$.

3.2. Remarks.

- It follows from the theorem that with probability one the results hold for a dense set of β . This implies that with probability one, (3.5)–(3.8) hold for all β . However, we have not shown whether or not for a particular realization, there might be an exceptional β for which (3.4) does not hold.
- The restriction to $t \in (0, 2]$ is only a convenience. By scaling we get a similar result for $t \in (0, \infty)$.
- The relationship $\dim_h[\gamma(\Theta_\beta)] = 2 \dim_h[\Theta_\beta]/(1 - \beta)$ can be understood as follows. For s small, the image of the interval $[t, t + s^2]$ under $\hat{f}_t$ can be approximated by a set of diameter $s|\hat{f}'_t(is)|$ containing $\hat{f}_t(is)$. If $|\hat{f}'_t(is)| \approx s^{-\beta}$, then this set has diameter $s^{1-\beta}$. That is to say, intervals of length (diameter) s^2 in a covering of Θ_β are sent to sets of diameter $s^{1-\beta}$. Note that this is in contrast to complex Brownian motion where intervals of length s^2 are always sent to sets whose diameter is of order s .
- Since $\Theta_\beta \subset \overline{\Theta}_\beta \cap \tilde{\Theta}_\beta \cap \underline{\Theta}_\beta^*$ and $\underline{\Theta}_\beta^* \subset \underline{\Theta}_\beta$, it suffices to prove the lower bounds for Θ_β in (3.4) and the upper bounds for $\overline{\Theta}_\beta, \underline{\Theta}_\beta, \underline{\Theta}_\beta^*$ in (3.5)–(3.8). The upper bounds will be proved in Section 4.1 and the lower bounds in Section 4.2.

- To prove the upper bound (3.5) it suffices to show for each $s > 0$,

$$\dim_h(\overline{\Theta}_\beta \cap (s, 2]) \leq \hat{d}_\beta,$$

and similarly for (3.6)–(3.8). This is what we do in Section 4.1.

- Recall that $\underline{\Theta}_\beta^* \subset \underline{\Theta}_\beta$. It is open whether or not

$$\dim_h[\gamma(\underline{\Theta}_\beta)] \leq d_\beta.$$

- Note that $(0, 2] = \overline{\Theta}_{\beta_*} \cup \underline{\Theta}_{\beta_*}^*$. It follows that

$$\dim_h(\gamma(0, 2]) = d_{\beta_*} = d.$$

Hence, Beffara's theorem on the dimension of the path $[1, 9]$ is a particular case of the theorem.

- The statements about the dimension of $\gamma(\Theta_\beta), \gamma(\overline{\Theta}_\beta), \gamma(\underline{\Theta}_\beta^*)$ are independent of the parametrization of the curve.
- Using the Markov property for SLE it is not hard to show that with probability one, either Θ_β is dense in $(0, \infty)$ or it is empty. Also, $\dim_h[\gamma(\Theta_\beta \cap [t_1, t_2])]$ is the same for all $0 < t_1 < t_2 \leq 2$. In particular, in order to prove the lower bound on dimension, it suffices to prove that for all $\alpha < d_\beta$,

$$\mathbb{P}\{\dim_h[\gamma(\Theta_\beta \cap [1, 2])] \geq \alpha\} > 0.$$

This is what we will do in Section 4.2. The proof proves the slightly stronger (for $\kappa > 4$) result

$$\mathbb{P}\{\dim_h[\mathbb{H} \cap \gamma(\Theta_\beta \cap [1, 2])] \geq \alpha\} > 0.$$

- If $\kappa = 8$, $\beta_* = \beta_+ = 1$ and $\dim_h[\gamma(\Theta_1)] = 2$. This is related to the fact that this is the hardest case to establish the existence of the curve; the curve is almost surely not Hölder continuous (in the capacity parameterization) when $\kappa = 8$ [5]. For other values of κ , $\beta_* < \beta_+ < 1$.

3.3. Multifractal spectrum of harmonic measure. Let $\Theta_\alpha^{\text{hm}}$ be defined as in Section 2.3. Let

$$F_{\text{tip}}(\alpha) := d_{1-1/\alpha} = \alpha \left(1 - \frac{4}{\kappa}\right) + \frac{(4 + \kappa)^2}{8\kappa} - \frac{\kappa}{8} \left(\frac{\alpha^2}{2\alpha - 1}\right),$$

and let $\alpha_-, \alpha_*, \alpha_+$ correspond to $\beta_-, \beta_*, \beta_+$ through the relation

$$\alpha = \frac{1}{1 - \beta}.$$

Remark. We can compare the function F_{tip} with the conjectured almost sure bulk spectrum for SLE_κ given by

$$F_{\text{bulk}}(\alpha) = \alpha + \frac{(4 + \kappa)^2}{8\kappa} - \frac{(4 + \kappa)^2}{8\kappa} \left(\frac{\alpha^2}{2\alpha - 1}\right).$$

Theorem 3.2. *Suppose $\alpha_- \leq \alpha \leq \alpha_+$. For chordal SLE_κ , with probability one,*

$$\dim_h[\gamma(\Theta_\alpha^{\text{hm}})] = F_{\text{tip}}(\alpha).$$

Proof. This is an immediate corollary of Theorem 3.1 and Lemma 2.9. $\square$

Theorem 3.2 combined with (2.20) gives some information on $\tilde{\Theta}_\alpha^{\text{hm}}$. In Section 6, we will use the forward Loewner flow to give a proof of the following.

Theorem 3.3. *If $0 < \kappa < 8$ and $1/2 \leq \alpha \leq \alpha_*$, then with probability one there exists a set V such that $\dim_h[\gamma(V)] \leq F_{\text{tip}}(\alpha)$ and for $t \notin V$, $\gamma(t) \in \mathbb{H}$,*

$$\tilde{\mu}(t, 2^{-n}) \asymp 2^{-n\alpha}, \quad n \rightarrow \infty. \quad (3.9)$$

Let

$$\tilde{T}_\alpha^{\text{hm}} = \{t \in (0, 2] : \tilde{\mu}(t, 2^{-n}) \gtrsim 2^{-n\alpha}, \gamma(t) \in \mathbb{H}\}$$

and note that Theorem 3.3 combined with (2.20) and Theorem 3.2 implies that for each $\alpha_- \leq \alpha < \alpha_*$ with probability one

$$\dim_h[\gamma(\tilde{T}_\alpha^{\text{hm}})] = F_{\text{tip}}(\alpha).$$

Indeed, it follows directly from (2.20) and Theorem 3.2 that the lower bound on the dimension holds with probability one. To get the upper bound, notice that $\tilde{T}_\alpha^{\text{hm}}$ is contained in $\{t \in (0, 2] : \tilde{\mu}(t, 2^{-n}) \gtrsim_{\text{i.o.}} 2^{-n\alpha}\}$, which, for those t such that $\gamma(t) \in \mathbb{H}$, in turn is contained in the set V from Theorem 3.3.

3.4. Parameters. In the statement of the main theorem, β and ρ were the parameters used. However, in deriving the result it is useful to consider a number of other parameters. Let

$$r_* = \min \left\{ 1, \frac{8}{\kappa} \right\}, \quad r_c = \frac{1}{2} + \frac{4}{\kappa},$$

and note that

$$0 < r_* \leq r_c,$$

where the second inequality is strict unless $\kappa = 8$. Let $r < r_c$; we define $\lambda, \zeta, \beta, \rho$ as functions of r .

Let

$$\lambda = \lambda(r) = r \left(1 + \frac{\kappa}{4} \right) - \frac{\kappa r^2}{8}. \quad (3.10)$$

We write $\lambda_* = \lambda(r_*)$, and similarly for other parameters. As r increases from $-\infty$ to r_c , λ increases from $-\infty$ to

$$\lambda_c = 1 + \frac{3\kappa}{32} + \frac{2}{\kappa}.$$

Since the relationship is injective, we can write either $\lambda(r)$ or $r(\lambda)$. Solving the quadratic equation gives

$$r(\lambda) = \frac{4 + \kappa - \sqrt{(4 + \kappa)^2 - 8\lambda\kappa}}{\kappa}.$$

Also,

$$\lambda(0) = 0, \quad \lambda_* = d.$$

Let

$$\zeta = \zeta(r) = r - \frac{\kappa r^2}{8} = \lambda(r) - \frac{\kappa r}{4}, \quad (3.11)$$

and note that

$$\zeta_* = 2 - d.$$

We can write ζ as a function of λ ,

$$\zeta(\lambda) = \lambda + \frac{\sqrt{(4 + \kappa)^2 - 8\lambda\kappa} - 4 - \kappa}{4}.$$

We now briefly discuss some results from [9, 5]. The reverse-time Loewner flow h_t (see Section 5.2 for definitions) has the property that for fixed t , the distribution of $|h'_t(z)|$ is the same as that of $|\hat{f}'_t(z)|$. For the reverse-time flow, if $r \in \mathbb{R}$ and λ, ζ are defined as above, then

$$|h'_t(z)|^\lambda Y_t(z)^\zeta [\sin \arg Z_t(z)]^{-r}$$

is a martingale. Typically one expects $Y_t(i) \asymp \sqrt{t}$ and $\sin \arg Z_t(i) \asymp 1$. If this is true, then the martingale property would imply

$$\mathbb{E} \left[|\hat{f}'_{t^2}(i)|^\lambda \right] = \mathbb{E} \left[|h'_{t^2}(i)|^\lambda \right] \asymp t^{-\zeta}.$$

It turns out that this argument can be carried out if $r < r_c$, and this is the starting point for determining the multifractal spectrum.

We define $\beta = \beta(r)$ by the relation

$$\frac{d\zeta}{d\lambda} = -\beta.$$

A straightforward calculation gives

$$\beta(r) = -1 + \frac{\kappa}{4 + \kappa - \kappa r}, \quad \kappa r(\beta) = 4 + \kappa - \frac{\kappa}{\beta + 1}.$$

Note that β increases with r with

$$\beta(-\infty) = -1, \quad \beta(0) = -\frac{4}{4 + \kappa} = \beta_\#, \quad \beta(r_*) = \beta_*, \quad \beta_c = 1,$$

where $\beta_\#, \beta_*$ are as defined in the previous section. Roughly speaking, $\mathbb{E}[|\hat{f}'_{t^2}(i)|^\lambda]$ is carried on an event on which $|\hat{f}'_{t^2}(i)| \approx t^\beta$ and

$$\mathbb{P} \left\{ |\hat{f}'_{t^2}(i)| \approx t^\beta \right\} \approx t^{-(\zeta + \lambda\beta)}. \quad (3.12)$$

We emphasize that the relation between r, λ, β for $-\infty < r < r_c$ is bijective and in order to specify the values of the parameters it suffices to give the value of any one of these. For example, we could choose β as the independent variable and write $r(\beta), \lambda(\beta)$. This is the natural approach when proving Theorem 3.1, but the formulas tend to be somewhat simpler if we choose r to be the independent variable.

From (3.12), it is natural to define

$$\rho = \rho(r) = \zeta(r) + \lambda(r) \beta(r) = \frac{\kappa^2 r^2}{8(4 + \kappa - \kappa r)}.$$

We can also write ρ as a function of β and a computation gives (3.1). Note that

$$\frac{d\rho}{d\beta} = \frac{d\zeta}{d\lambda} \frac{d\lambda}{d\beta} + \lambda + \beta \frac{d\lambda}{d\beta} = \lambda.$$

Let r_+, r_- denote the two values of r for which $\zeta(r) + \lambda(r) \beta(r) = 2$ with corresponding values $\beta_+ = \beta(r_+), \beta_- = \beta(r_-)$. Then

$$r_{\pm} = \frac{4}{\kappa} [-2 \pm \sqrt{8 + \kappa}],$$

and β_+, β_- are given as in (3.2)–(3.3). Note that if $\kappa \neq 8$, then $r_+ < r_c$.

Define

$$d_{\beta} = \frac{2 - \rho(\beta)}{1 - \beta} = \frac{2 - (\zeta + \beta\lambda)}{1 - \beta}.$$

Note that d_{β} is maximized at $\beta = \beta_*$ (interpreted as a limit for $\kappa = 8$) with $d_* = d$. We can also define d as a function of r ,

$$d(r) = 1 + \frac{\kappa - \frac{\kappa^2 r^2}{8}}{8 + \kappa - 2\kappa r}.$$

Straightforward differentiation shows that $d'(r) = 0$ implies $r = 1$ or $r = 8/\kappa$. Note that $1 = 8/\kappa = r_+$ if $\kappa = 8$ and

$$1 < r_+ < \frac{8}{\kappa}, \quad \kappa < 8,$$

$$\frac{8}{\kappa} < r_+ < 1, \quad \kappa > 8.$$

From this we can see that $d(r)$ achieves its maximum on $(-\infty, r_+)$ at $r = r_*$; in fact, $d(\beta)$ increases for $\beta < \beta_*$ and decreases for $\beta_* < \beta < \beta_+$.

In order to match the notation of [9], let

$$q = r_c - r = \frac{1}{2} + \frac{4}{\kappa} - r. \quad (3.13)$$

Obviously, $q > 0$ if and only if $r < r_c$. For future reference we note that

$$\frac{1 - 2q}{1 + 2q} = \beta. \quad (3.14)$$

4. PROOF OF THE MAIN THEOREM

In this section we will present the proof of the theorem relying on estimates about moments of derivatives of the map $\hat{f}$. The upper bounds are proved in Section 4.1, and the lower bound is proved in Section 4.2.

4.1. Upper bounds. In this subsection (and this subsection only) we write

$$\hat{f}_{j,n} = \hat{f}_{(j-1)2^{-2n}}.$$

For each $t \in [0, \infty)$, we associate a dyadic time by defining

$$t_n = t_n(t) = \frac{j-1}{2^{2n}} \quad \text{if} \quad \frac{j-1}{2^{2n}} \leq t < \frac{j}{2^{2n}}.$$

We fix s with $0 < s \leq 2$ and allow constants to depend on s .

The next theorem states the derivative estimates that we will use for the upper bounds; a proof can be found in [5].

Theorem 4.1. [5] *If $r < r_c$, there exists $c < \infty$ such that for all $t \geq 1$,*

$$\mathbb{E} \left[|\hat{f}'_{t^2}(i)|^\lambda \right] \leq c t^{-\zeta}. \quad (4.1)$$

Corollary 4.2. *If $\beta \geq \beta_\#$, there is a $c < \infty$ such that if*

$$N_{n,\beta} = \sum_{s2^{2n} \leq j \leq 2^{2n+1}} 1 \left\{ |\hat{f}'_{j,n}(i2^{-n})| \geq 2^{n\beta} \right\},$$

then

$$\mathbb{E}[N_{n,\beta}] \leq c 2^{n(2-\rho)}. \quad (4.2)$$

Proof. The range $\beta \geq \beta_\#$ corresponds to $\lambda \geq 0$. Hence, by Chebyshev's inequality,

$$\begin{aligned} \mathbb{P}\{|\hat{f}'_{j,n}(i2^{-n})| \geq 2^{n\beta}\} &\leq 2^{-n\beta\lambda} \mathbb{E} \left[|\hat{f}'_{j,n}(i2^{-n})|^\lambda \right] \\ &= 2^{-n\beta\lambda} \mathbb{E} \left[|\hat{f}'_j(i)|^\lambda \right] \leq c 2^{-n\beta\lambda} j^{-\zeta/2}, \end{aligned}$$

and hence

$$\begin{aligned} \mathbb{E}[N_{n,\beta}] &= \sum_{s2^{2n} \leq j \leq 2^{2n+1}} \mathbb{P}\{|\hat{f}'_{j,n}(i2^{-n})| \geq 2^{n\beta}\} \\ &\leq 2^{-n\beta\lambda} \sum_{s2^{2n} \leq j \leq 2^{2n+1}} j^{-\zeta/2} \\ &\leq c 2^{n(2-\beta\lambda-\zeta)} = c 2^{n(2-\rho)}. \end{aligned}$$

□

Corollary 4.3. *If $\beta < \beta_\#$, there is a $c < \infty$ such that if*

$$N_{n,\beta}^* = \sum_{s2^{2n} \leq j \leq 2^{2n+1}} 1 \left\{ |\hat{f}'_{j,n}(i2^{-n})| \leq 2^{n\beta} \right\},$$

then

$$\mathbb{E}[N_{n,\beta}^*] \leq c 2^{n(2-\rho)}. \quad (4.3)$$

Proof. This is proved in the same way using $\lambda < 0$. □

The standard technique to find upper bounds for Hausdorff dimension uses an appropriate sequences of covers for a set. We will now describe the covers that we will use. Let

$$I(j, n) = \left[\frac{j-1}{2^{2n}}, \frac{j}{2^{2n}} \right].$$

If $b, \bar{b} \in \mathbb{R}$ with $-1 < b < \bar{b} < 1$, let $\bar{\mathcal{B}}(j, n, \bar{b})$ be the closed disk in $\mathbb{C}$ of radius $2^{n(\bar{b}-1)}$ centered at $\hat{f}_{j,n}(i2^{-n})$, and let

$$I_n(s, b) = \bigcup I(j, n) \quad \mathcal{B}_n(s, b, \bar{b}) = \bigcup \bar{\mathcal{B}}(j, n, \bar{b}),$$

where in each case the union is over $s2^{2n} \leq j \leq 2^{2n+1}$ with $|\hat{f}'_{j,n}(i2^{-n})| \geq 2^{nb}$. Let

$$I^m(s, b) = \bigcup_{n=m}^{\infty} I_n(s, b), \quad \mathcal{B}^m(s, b, \bar{b}) = \bigcup_{n=m}^{\infty} \mathcal{B}_n(s, b, \bar{b}).$$

Lemma 4.4. *If $-1 < b < \beta < b_1 < \bar{b} < 1$, then for each m ,*

$$\bar{\Theta}_\beta \cap (s, 2] \subset I^m(s, b),$$

$$\gamma \left(\bar{\Theta}_\beta \cap \tilde{\Theta}_{b_1} \cap (s, 2] \right) \subset \mathcal{B}^m(s, b, \bar{b}).$$

Proof. Suppose $t \in \bar{\Theta}_\beta \cap (s, 2]$. By Proposition 2.8, there exists a subsequence $n_j \rightarrow \infty$ such that

$$|f'_{t_{n_j}}(i2^{-n_j})| \geq 2^{n_j b}.$$

In other words, there is a sequence n_j such that $I(t_{n_j} 2^{2n_j}, n_j) \in I_{n_j}(s, b)$. This proves the first assertion.

If $t \in \bar{\Theta}_\beta \cap \tilde{\Theta}_{b_1} \cap (s, 2]$ and $b_1 < u < \bar{b}$, then (2.19) shows that for all n sufficiently large,

$$|\gamma(t) - \hat{f}_t(i2^{-n})| \leq 2^{(u-1)n}.$$

The triangle inequality gives

$$|\gamma(t) - \hat{f}_{t_n}(i2^{-n})| \leq |\gamma(t) - \hat{f}_t(i2^{-n})| + |\hat{f}_t(i2^{-n}) - \hat{f}_{t_n}(i2^{-n})|,$$

and estimating as in (2.17), we have for n sufficiently large

$$|\hat{f}_t(i2^{-n}) - \hat{f}_{t_n}(i2^{-n})| \leq 2^{(u-1)n}.$$

Hence, for n sufficiently large,

$$|\gamma(t) - \hat{f}_{t_n}(i2^{-n})| \leq 2^{n_j(\bar{b}-1)}.$$

This implies for all j sufficiently large,

$$\gamma(t) \in \mathcal{B}(t_{n_j} 2^{2n_j}, b, \bar{b}).$$

□

Proposition 4.5. *If $\beta \geq \beta_\#$, then with probability one,*

$$\dim_h(\overline{\Theta}_\beta \cap (s, 2]) \leq \hat{d}_\beta, \quad (4.4)$$

Moreover, if $\beta > \beta_+$, then with probability one

$$\overline{\Theta}_\beta \cap (s, 2] = \emptyset.$$

Proof. It suffices to consider $\beta_\# < \beta < 1$. Suppose $\beta_\# < b < \beta < 1$. Using the cover from Lemma 4.4, we get

$$\mathcal{H}^\alpha [\overline{\Theta}_\beta \cap (s, 2)] \leq \sum_{n=m}^{\infty} N_{n,b} 2^{-2\alpha n},$$

and hence (4.2) implies

$$\mathbb{E}(\mathcal{H}^\alpha [\overline{\Theta}_\beta \cap (s, 2)]) \leq c \sum_{n=m}^{\infty} 2^{n(2-\rho(b))} 2^{-2\alpha n}.$$

The sum goes to zero, provided that $2\alpha > 2 - \rho(b)$, and hence with probability one

$$\mathcal{H}^\alpha(\overline{\Theta}_\beta \cap (s, 2]) = 0, \quad \alpha > \frac{2 - \rho(b)}{2}.$$

Letting $b \rightarrow \beta$ gives (4.4).

For the second assertion, note that

$$\mathbb{P}\{\overline{\Theta}_\beta \cap (s, 2] \neq \emptyset\} \leq \sum_{n=m}^{\infty} \mathbb{E}[N_{n,b}] \leq c \sum_{n=m}^{\infty} 2^{n(2-\rho(b))}.$$

If $\beta > \beta_+$, then $\rho(\beta) > 2$ and we can find $b < \beta$ with $\rho(b) > 2$. □

Lemma 4.6. *If $\beta_\# \leq \beta < b_1 < 1$, then with probability one,*

$$\dim_h [\gamma(\overline{\Theta}_\beta \cap \tilde{\Theta}_{b_1} \cap (s, 2])] \leq \frac{2 - \rho(\beta)}{1 - b_1}. \quad (4.5)$$

Proof. Choose $b, \bar{b}$ with $\beta_\# < b < \beta < b_1 < \bar{b} < 1$. Using the cover from Lemma 4.4, we get

$$\mathcal{H}^\alpha [\gamma(\overline{\Theta}_\beta \cap \tilde{\Theta}_{b_1} \cap (s, 2])] \leq \sum_{n=m}^{\infty} N_{n,b} 2^{\alpha(\bar{b}-1)n},$$

and hence (4.2) implies

$$\mathbb{E}(\mathcal{H}^\alpha [\gamma(\overline{\Theta}_\beta \cap \tilde{\Theta}_{b_1} \cap (s, 2])]) \leq c \sum_{n=m}^{\infty} 2^{n(2-\rho(b))} 2^{\alpha(\bar{b}-1)n}.$$

The sums on the right go to zero provided that

$$\alpha > \frac{2 - \rho(b)}{1 - \bar{b}},$$

respectively. We now choose a sequence of values for $b, \bar{b}$ that converge to β, b_1 to conclude (4.5). □

Proposition 4.7. *If $\beta_* \leq \beta \leq 1$, then with probability one,*

$$\dim_h [\gamma(\overline{\Theta}_\beta \cap (s, 2])] \leq d_\beta. \quad (4.6)$$

Proof. If $\beta = b_0 < b_1 < b_2 < \dots < b_k < 1$ with $b_k > b_+$, then

$$\overline{\Theta}_\beta = \bigcup_{j=1}^k (\overline{\Theta}_{b_{j-1}} \cap \tilde{\Theta}_{b_j}).$$

Therefore, (4.5) implies

$$\dim_h [\gamma(\overline{\Theta}_\beta \cap (s, 2])] \leq \max \left\{ \frac{2 - \rho(b_{j-1})}{1 - b_j} : j = 1, \dots, k \right\}.$$

By taking finer and finer partitions and using the continuity of ρ , we see that

$$\dim_h [\gamma(\overline{\Theta}_\beta \cap (s, 2])] \leq \sup_{b \geq \beta} \frac{2 - \rho(b)}{1 - b} = d_\beta.$$

The last equality uses $\beta \geq \beta_*$ and the fact which can easily be verified (see Section 3.4) that the function

$$b \mapsto \frac{2 - \rho(b)}{1 - b},$$

is decreasing on the interval $[\beta_*, \beta_+]$. $\square$

For $\beta < \beta_\#$ we use a slightly different cover. Let $I(j, n)$ be as above and

$$I_n^*(s, b) = \bigcup I(j, n) \quad \mathcal{B}_n^*(s, b, \bar{b}) = \bigcup \mathcal{B}(j, n, \bar{b}),$$

where in each case the union is over $s2^{2n} \leq j \leq 2^{2n+1}$ with $|\hat{f}'_{j,n}(i2^{-n})| \leq 2^{nb}$. Let

$$I_*^m(s, b) = \bigcup_{n=m}^{\infty} I_n^*(s, b), \quad \mathcal{B}_*^m(s, b, \bar{b}) = \bigcup_{n=m}^{\infty} \mathcal{B}_n^*(s, b, \bar{b}).$$

Lemma 4.8. *If $-1 < \beta < b < \bar{b}$,*

$$\begin{aligned} \underline{\Theta}_\beta \cap (s, 2] &\subset I_*^m(s, b), \\ \gamma(\underline{\Theta}_\beta^* \cap (s, 2]) &\subset \mathcal{B}_*^m(s, b, \bar{b}). \end{aligned}$$

Proof. Suppose $t \in \underline{\Theta}_\beta \cap (s, 2]$. By Proposition 2.8, there exists a subsequence $n_j \rightarrow \infty$ such that

$$|\hat{f}'_{t_{n_j}}(i2^{-n_j})| \leq 2^{n_j b}. \quad (4.7)$$

In other words, there is a sequence n_j such that $I(t_{n_j} 2^{2n_j}, n_j) \in I_{n_j}^*(s, b)$. This proves the first assertion.

Now suppose $t \in \underline{\Theta}_\beta^* \cap (s, 2] \subset \underline{\Theta}_\beta \cap (s, 2]$. Then there exists a sequence n_j such that both (4.7) holds and

$$v_t(2^{-n_j}) \leq 2^{n_j(b-1)}.$$

Using the triangle inequality as in Lemma 4.4, we see that

$$|\gamma(t) - \hat{f}_{t_{n_j}}(i2^{-n_j})| \leq v_t(2^{-n_j}) + \left| \hat{f}_t(i2^{-n_j}) - \hat{f}_{t_{n_j}}(i2^{-n_j}) \right|,$$

and arguing as before we see that for j sufficiently large

$$|\gamma(t) - \hat{f}_{t_{n_j}}(i2^{-n_j})| \leq 2^{n_j(\bar{b}-1)},$$

and

$$\gamma(t) \in \mathcal{B}^*(t_{n_j}2^{2n}, b, \bar{b}).$$

□

Proposition 4.9. *If $\beta < \beta_\#$, then with probability one,*

$$\dim_h(\underline{\Theta}_\beta \cap (s, 2]) \leq \hat{d}_\beta,$$

$$\dim_h(\gamma(\underline{\Theta}_\beta^* \cap (s, 2])) \leq d_\beta.$$

Moreover, if $\beta < \beta_-$, with probability one,

$$\underline{\Theta}_\beta \cap (s, 2] = \emptyset.$$

Proof. This is proved in the same way as Proposition 4.5 and Lemma 4.6 using (4.3). □

4.2. Lower bound. In this subsection we prove the lower bound for the dimension in (3.4). We fix r such that $\rho = \lambda\beta + \zeta < 2$ and recall that $r < r_c$. As has been pointed out, it suffices to show that with positive probability

$$\dim_h(\Theta_\beta \cap [1, 2]) \geq \frac{2-\rho}{2}, \quad \dim_h[\gamma(\Theta_\beta \cap [1, 2])] \geq \frac{2-\rho}{1-\beta}. \quad (4.8)$$

We will use a standard technique of Frostman to show that with positive probability there exist nontrivial positive measures μ, ν whose supports are contained in $\Theta_\beta \cap [1, 2]$ and $\gamma(\Theta_\beta \cap [1, 2])$, respectively, such that

$$\mathcal{E}_\alpha(\mu) < \infty, \quad \alpha < \frac{2-\rho}{2}; \quad \mathcal{E}_\alpha(\nu) < \infty, \quad \alpha < \frac{2-\rho}{1-\beta},$$

where

$$\mathcal{E}_\alpha(\mu) = \int \int \frac{\mu(dx) \mu(dy)}{|x-y|^\alpha}$$

is the energy integral. It is well known that this implies (4.8). For this subsection, we will adopt a different notation than in the previous subsection. We let

$$\hat{f}_{j,n} = 1 + \frac{(j-1)}{n^2}, \quad j = 1, 2, \dots, n^2.$$

We will be studying $|\hat{f}'_{j,n}(i/n)|$. The proof considers a subset of times in $\Theta_\beta \cap [1, 2]$ that behave in some sense nicely. The hard work is Theorem 4.10 which will be proved in Section 5. This theorem discusses the existence of some events $E_{j,n}$ on which

$$|\hat{f}'_{j,n}(iy)| \approx y^{-\beta}, \quad n^{-1} \leq y \leq 1.$$

The definition of the events (“good times”) will be left for Section 5.

Theorem 4.10. *Suppose $\rho = \lambda\beta + \zeta < 2$. There exist $c < \infty$, a subpower function ψ , and events*

$$E_{j,n}, \quad n = 1, 2, \dots, \quad j = 1, \dots, n^2,$$

such that the following hold. Let $E(j, n) = 1_{E_{j,n}}$ and

$$F(j, n) = n^{\zeta-2} |\hat{f}'_{j,n}(i/n)|^\lambda E(j, n).$$

- *If $1 \leq j \leq n^2$, then on the event $E_{j,n}$,*

$$\psi(1/y)^{-1} y^{-\beta} \leq |\hat{f}'_{j,n}(iy)| \leq \psi(1/y) y^{-\beta}, \quad n^{-1} \leq y \leq 1. \quad (4.9)$$

- *If $1 \leq j \leq k \leq n^2$,*

$$c_1 n^{-2} \leq \mathbb{E}[F(j, n)] \leq n^{\zeta-2} \mathbb{E} \left[|\hat{f}'_{j,n}(i/n)|^\lambda \right] \leq c_2 n^{-2}, \quad (4.10)$$

- *If $1 \leq j \leq k \leq n^2$,*

$$\mathbb{E}[F(j, n) F(k, n)] \leq n^{-4} \left(\frac{n^2}{k-j+1} \right)^{\frac{\rho}{2}} \psi \left(\frac{n^2}{k-j+1} \right). \quad (4.11)$$

- *If $1 \leq j < k \leq n^2$ and $E(j, n) E(k, n) = 1$,*

$$\left| \hat{f}_{j,n}(i/n) - \hat{f}_{k,n}(i/n) \right|^{-\frac{2-\rho}{1-\beta}} \leq \left(\frac{n^2}{k-j+1} \right)^{\frac{2-\rho}{2}} \psi \left(\frac{n^2}{k-j+1} \right)^{d_\beta}. \quad (4.12)$$

Proof. This theorem combines Propositions 5.8 and 5.9 proved in Section 5. $\square$

Proposition 4.11. *Under the assumptions of Theorem 4.10, with positive probability there exists $A \subset [1, 2]$ such that for $t \in A$,*

$$\frac{1}{4} y^{-\beta} \psi(1/y)^{-1} \leq |\hat{f}'_t(iy)| \leq 4 y^{-\beta} \psi(1/y), \quad 0 < y \leq 1. \quad (4.13)$$

and such that

$$\dim_h(A) \geq \frac{2-\rho}{2}, \quad \dim_h[\gamma(A)] \geq \frac{2-\rho}{1-\beta}.$$

Proof assuming Theorem 4.10. We use a now standard argument to show that with positive probability a “Frostman measure” of appropriate dimension can be put on the set of t satisfying (4.13). The proof is very similar to that of [9, Lemma 10.3] so we omit some of the details.

Let $\mu_{j,n}$ denote the random measure on $\mathbb{R}$ that is a multiple of Lebesgue measure on $I(j, n) := [1 + (j-1)n^{-2}, 1 + jn^{-2}]$ where the multiple is chosen so that $\|\mu_{j,n}\| = F(j, n)$. Let $\nu_{j,n}$ denote the random measure on $\mathbb{C}$ that is a multiple of Lebesgue measure on the disk of radius $n^{\beta-1} \psi(n^2)^{-1}/4$ centered at $\hat{f}_{j,n}(i/n)$ where the constant is chosen so that $\|\nu_{j,n}\| = F(j, n)$. Let $\mu_n = \sum_{j=1}^{n^2} \mu_{j,n}$, $\nu_n = \sum_{j=1}^{n^2} \nu_{j,n}$. Note that

$$\|\mu_n\| = \|\nu_n\| = \sum_{j=1}^{n^2} F(j, n).$$

From (4.10) and (4.11), we see that

$$\mathbb{E} [\|\mu_n\|] \geq c_1, \quad \mathbb{E} [\|\mu_n\|^2] \leq c_2.$$

Hence

$$\mathbb{P} \{\|\mu_n\| > 0\} \geq c > 0,$$

uniformly in n . From (4.11) and (4.12), we can show that there is a C_α such that

$$\mathbb{E}[\mathcal{E}_\alpha(\mu_n)] \leq C_\alpha, \quad \alpha < \frac{2-\rho}{2},$$

$$\mathbb{E}[\mathcal{E}_\alpha(\nu_n)] \leq C_\alpha, \quad \alpha < \frac{2-\rho}{1-\beta}.$$

We let μ denote a subsequential limit of the μ_n which with positive probability we know is nontrivial and satisfies $\mathcal{E}_\alpha(\mu) < \infty$ for all $\alpha < (2-\rho)/2$. Hence,

$$\dim_h[\text{supp } \mu] \geq \frac{2-\rho}{2}.$$

Similarly, let ν denote a subsequential limit of the ν_n which is nonzero with positive probability and satisfies

$$\dim_h[\text{supp } \nu] \geq \frac{2-\rho}{1-\beta}.$$

We claim that every $t \in \text{supp } \mu$ satisfies (4.13). Indeed, the construction shows that if $t \in \text{supp } \mu$, then there is a subsequence $n_k \rightarrow \infty$ and $j_k \in \{1, \dots, n_k^2\}$ such that $E(j_k, n_k) = 1$ and

$$\lim_{k \rightarrow \infty} \frac{j_k - 1}{n_k^2} = t. \tag{4.14}$$

Suppose for some $t \in [1, 2]$ and $0 < y \leq 1$, we had

$$|\hat{f}'_t(iy)| \geq 4y^{-\beta} \psi(1/y).$$

Continuity would imply that for all s in a neighborhood of t ,

$$|\hat{f}'_s(iy)| \geq 2y^{-\beta} \psi(1/y).$$

This implies that there is no sequence (j_k, n_k) as above with $E(j_k, n_k)$ satisfying (4.14). A similar argument shows that there cannot exist $t \in \text{supp } \mu$ and y with $|\hat{f}'_t(iy)| \leq (1/4)y^{-\beta} \psi(1/y)^{-1}$, and this gives (4.13). Similarly, $\text{supp } \nu$ is contained in $\gamma(A')$, where A' denotes the set of $t \in [1, 2]$ satisfying (4.13). $\square$

5. ESTIMATING THE MOMENTS

In this section $r < r_c$ with corresponding values of $\zeta, \beta, \rho, \lambda$. All constants may depend on r . Let us give an overview of the section. We begin by discussing the reverse-time Loewner flow and how it relates to f_t . We then go on to define the “good times” which, roughly speaking, are T for which the reverse flow driven by $t \mapsto V_{T-t} - V_t$ behaves in some sense nicely. (Here, V_t is a two-sided Brownian motion.) We make this precise in a number of lemmas that show how $|h'_t| = |h'_{T,t}|$ can be controlled on the event that T is “good”. The needed correlation estimates can then be derived using moment bounds from [9, 5].

5.1. Reverse Loewner flow. Here we state the basic lemma that relates the reverse Loewner flow to the forward flow for SLE. We will estimate the moments for $h, \tilde{h}$ rather than for $\hat{f}$.

If V_t is a continuous function, define g_t to be the solution to the forward-time (chordal) Loewner equation

$$\partial_t g_t(z) = \frac{a}{g_t(z) - V_t}, \quad g_0(z) = z. \quad (5.1)$$

Let $f_t(z) = g_t^{-1}(z)$, $\hat{f}_t(z) = f_t(z + V_t)$. If U_t is another continuous function, let h_t be the solution to the reverse-time (chordal) Loewner equation

$$\partial_t h_t(z) = \frac{a}{U_t - h_t(z)}, \quad h_0(z) = z. \quad (5.2)$$

The next lemma relates the forward-time and reverse-time equations; although versions of this have appeared before we give a short proof. We point out that this is a fact about the Loewner equation itself; no assumptions are made about the function V other than continuity.

Lemma 5.1. *Assume $V_t, -\infty < t < \infty$, is a continuous function with $V_0 = 0$. For each $T \geq 0$, let*

$$U_{t,T} = V_{T-t} - V_T.$$

Let $g_t, 0 \leq t < \infty$, be the solution to the forward-time Loewner equation (5.1), and let $f_t, \hat{f}_t$ be as above. Let $h_{t,T}, 0 \leq t < \infty$, be the solution to the reverse-time Loewner equation (5.2) with $U_t = U_{t,T}$. Let

$$Z_{t,T}(z) = h_{t,T}(z) - U_{t,T}.$$

Then

$$h_{T,T}(z) = \hat{f}_T(z) - V_T = \hat{f}_T(z) + U_T,$$

and if $0 \leq S \leq T$, $z, w \in \mathbb{H}$,

$$h_{T,T}(z) = h_{S,S}(Z_{T-S,T}(z)) + U_{T-S,T}, \quad (5.3)$$

$$\hat{f}_T(z) - \hat{f}_S(w) = h_{S,S}(Z_{T-S,T}(z)) - h_{S,S}(w). \quad (5.4)$$

In particular,

$$h'_{T,T}(z) = h'_{S,S}(Z_{T-S,T}(z)) h'_{T-S,T}(z).$$

Proof. Fix T and let $U_t = U_{t,T}$, $h_t = h_{t,T}$. For $0 \leq s \leq T$, we have

$$U_{t,s} = V_{s-t} - V_s = U_{T-s+t} - U_{T-s}.$$

Let $u_t(z) = h_{T-t}(z) + V_T$. Then u_t satisfies

$$\partial_t u_t(z) = -\partial_t h_{T-t}(z) = \frac{a}{h_{T-t}(z) - [V_t - V_T]} = \frac{a}{u_t(z) - V_t}.$$

Also, $u_T(z) = h_0(z) + V_T = z + V_T$. By comparison, with (5.1),

$$u_0(z) = g_t^{-1}(z + V_T) = \hat{f}_t(z),$$

and hence

$$h_T(z) = u_0(z) - V_T = \hat{f}_T(z) - V_T = \hat{f}_T(z) + U_T.$$

For $0 \leq s \leq T$, define $h_t^{(s)}$ by

$$h_{s+t} = h_t^{(s)} \circ h_s.$$

By (5.2) we see that

$$\begin{aligned} \partial_t [h_t^{(s)}(z) - U_s] &= \frac{a}{U_{t+s} - h_t^{(s)}(z)} = \frac{a}{U_{t+s} - U_s - [h_t^{(s)}(z) - U_s]}, \\ h_0^{(s)}(z) - U_s &= z - U_s. \end{aligned}$$

Therefore,

$$h_t^{(s)}(z) - U_s = h_{t,T-s}(z - U_s), \quad (5.5)$$

which implies

$$h_{s+t}(z) = h_{t,T-s}(Z_{s,T}(z)) + U_s. \quad (5.6)$$

Setting $s = T - S$ and $t = S$ gives (5.3) and setting $s = S$, $t = T - S$ gives

$$\begin{aligned} \hat{f}_{T-S}(w) &= h_{T-S}(w) - U_{T-S,T-S} = h_{T-S,T-S}(w) - [U_T - U_S], \\ \hat{f}_T(w) &= h_T(z) - U_T = h_{T-S,T-S}(Z_{S,T}(z)) + U_S - U_T. \end{aligned}$$

Subtracting these equations gives (5.4). The final assertion follows from (5.3) and the chain rule. $\square$

The preceding lemma holds for all continuous V_t . If V_t is a standard Brownian motion, then so is $U_{t,T}$ for each T . We get the following corollary.

Lemma 5.2. *Suppose $0 < S < T$ and g_t , $0 \leq t \leq T$, is the solution to (5.1) where V_t is a standard Brownian motion. Suppose U_t is a standard Brownian motion and $h, \tilde{h}$ are the solutions to*

$$\begin{aligned} \partial_t h_t(z) &= \frac{a}{U_t - h_t(z)}, \quad h_0(z) = z, \\ \partial_t \tilde{h}_t(z) &= \frac{a}{\tilde{U}_t - \tilde{h}_t(z)}, \quad \tilde{h}_0(z) = z, \end{aligned} \quad (5.7)$$

where $\tilde{U}_t = U_{T-S+t} - U_{T-S}$. Then

$$h_{T-S+t}(z) = \tilde{h}_t(h_{T-S}(z) - U_{T-S}) + U_{T-S}.$$

Moreover, the joint distribution of the functions

$$\left(\hat{f}'_S(w), \operatorname{Im} \hat{f}_S(w), \hat{f}'_T(z), \operatorname{Im} \hat{f}_T(w), \hat{f}_T(z) - \hat{f}_S(w) \right)$$

is the same as the joint distribution of

$$\left(\tilde{h}'_S(w), \operatorname{Im} \tilde{h}_S(w), \tilde{h}'_S(Z), h'_{T-S}(z), \operatorname{Im} \tilde{h}_S(Z), \tilde{h}_S(Z) - \tilde{h}_S(w) \right),$$

where $Z = Z_{T-S}(z) = h_{T-S}(z) - U_{T-S}$.

5.2. Good times. Suppose that $T > 0$ and $h_t = h_{t,T}$ is defined as in the proof of Lemma 5.1. More specifically, g_t is the solution to the forward-time Loewner equation (5.1) with a (two-sided) Brownian motion V_t as driving function, and h_t is the solution to the reverse-time Loewner equation (5.2) with $U_t = U_{t,T} = V_{T-t} - V_T$ as driving function. Let

$$Z_t(z) = X_t(z) + Y_t(z) = h_t(z) - U_t.$$

Recall from Lemma 5.1 that we have

$$h_{t+s}(z) = h_{t+s,T}(z) = h_{t,T-S}(Z_s(z)) + U_s;$$

Note that

$$h'_{t+s}(z) = h'_{t,T-S}(Z_s(z)) h'_s(z). \quad (5.8)$$

If ψ is a subpower function and $0 < \delta \leq 1$, we let

$$\hat{\psi}_\delta(t) = \min\{\psi(t/\delta), \psi(1/t)\} = \begin{cases} \psi(t/\delta) & \text{if } t \leq \sqrt{\delta} \\ \psi(1/t) & \text{if } t > \sqrt{\delta} \end{cases}.$$

Note that for every subpower function ψ and every $c < \infty$, there is an $M < \infty$ such that for all $\delta > 0$,

$$\hat{\psi}_\delta(t) \leq M \quad \text{if } \delta \leq t \leq c\delta \text{ or } t \geq 1/c. \quad (5.9)$$

Roughly speaking, $\hat{\psi}_\delta(t)$ is $O(1)$ for t comparable to δ or comparable to 1 but can be larger for other $\delta < t < 1$.

Definition. We call a time T ψ -good at δ if the following five conditions hold for $h_t = h_{t,T}$ with $\hat{\psi} = \hat{\psi}_\delta$ and $Z_t = X_t + iY_t = Z_t(\delta i)$.

$$Y_{t^2} \geq t \hat{\psi}(t)^{-1}, \quad \delta \leq t \leq 2, \quad (5.10)$$

$$|X_{t^2}| \leq [t + \delta] \hat{\psi}(t), \quad 0 \leq t \leq 2, \quad (5.11)$$

$$(t/\delta)^\beta \hat{\psi}(t)^{-1} \leq |h'_{t^2}(i\delta)| \leq (t/\delta)^\beta \hat{\psi}(t), \quad \delta \leq t \leq 2, \quad (5.12)$$

$$t^{-\beta} \hat{\psi}(t)^{-1} \leq \frac{|h'_4(i\delta)|}{|h'_{t^2}(i\delta)|} = |h'_{4-t^2,T-t^2}(Z_{t^2})| \leq t^{-\beta} \hat{\psi}(t), \quad \delta \leq t \leq 2, \quad (5.13)$$

This definition depends on ψ and δ . Note that if T is ψ -good at δ and ϕ is a subpower function with $\psi \leq \phi$, then $\psi_\delta \leq \phi_\delta$ and T is ϕ -good at δ . In the remainder of this subsection, we derive some properties of ψ -good times. These will be used in the later subsections to estimate first and second moments for $|h'_{t^2}(\delta i)|^\lambda$ on the event that T is ψ -good at δ .

Proposition 5.3. *For every subpower function ψ there is a subpower function ϕ such that for all $\delta > 0$, if T is ψ -good at δ , then*

$$|U_{t^2}| \leq t\phi(1/t), \quad \delta \leq t \leq 2. \quad (5.14)$$

Proof. Let $X_t = X_t(i\delta)$, $Y_t = Y_t(i\delta)$. We let ϕ denote a subpower function whose value may change from line to line. From the Loewner equation, we know that

$$dX_t = -\frac{aX_t}{X_t^2 + Y_t^2} dt - dU_t.$$

Hence,

$$|U_{t^2}| \leq |X_{t^2}| + a \int_0^{t^2} \frac{|X_s| ds}{X_s^2 + Y_s^2}.$$

By (5.11), it suffices to show that

$$\int_0^{t^2} \frac{|X_s| ds}{X_s^2 + Y_s^2} \leq \phi(1/t).$$

Using (5.10) and (5.11), we have

$$\frac{|X_s|}{X_s^2 + Y_s^2} \leq \frac{\phi(1/s)}{\sqrt{s}},$$

and hence

$$\int_0^{t^2} \frac{|X_s| ds}{X_s^2 + Y_s^2} \leq \int_0^{t^2} \frac{\phi(1/s) ds}{\sqrt{s}} = \int_{t^{-2}}^\infty \phi(x) x^{-3/2} dx = \tilde{\phi}(1/t) t,$$

where

$$\tilde{\phi}(1/t) = t^{-1} \int_0^{t^2} \frac{\phi(1/s) ds}{\sqrt{s}}.$$

It is easy to check that $\tilde{\phi}$ is continuous and decays faster than x^ϵ for each ϵ . $\square$

Lemma 5.4. *If ψ is a subpower function, there is a $c > 0$ such that for every $0 < \delta \leq 1$, if T is ψ -good at δ and $Y_t = Y_t(\delta i)$, then*

$$Y_{\delta^2} \geq (1+c)\delta. \quad (5.15)$$

Proof. Using (5.10), (5.11) and the fact that Y_{t^2} increases with t , we see there is a $c_1 < \infty$ such that

$$|X_{t^2}| \leq c_1 \delta \leq c_1 Y_{t^2}, \quad 0 \leq t \leq \delta.$$

The Loewner equation (5.2) implies that

$$\partial_s Y_s = \frac{aY_s}{X_s^2 + Y_s^2} \geq \frac{a}{c_1^2 + 1} \frac{1}{Y_s}, \quad s \leq \delta^2,$$

from which (5.15) follows. $\square$

Lemma 5.5. *For every subpower function ψ , there is a c such that if $0 < \delta \leq 1$ and T is ψ -good at δ , then*

$$|h_{4-t^2, T-t^2}(Z_{t^2}) - h_{4-t^2, T-t^2}(\delta i)| \geq c \hat{\psi}_\delta(t)^{-2} t^{1-\beta}, \quad \delta \leq t \leq 2.$$

Proof. Using (5.10) and (5.15), we see that there is a $c_1 > 0$ such that if $\mathcal{B}$ denotes the open disk of radius $c_1 t \hat{\psi}_\delta(t)^{-1}$ about Z_{t^2} , then $\delta i \notin \mathcal{B}$. Using (5.13) and the Koebe (1/4)-theorem we see that $h_{4-t^2, T-t^2}(\mathcal{B})$ contains a disk of radius $(c_1/4) \hat{\psi}_\delta(t)^{-2} t^{1-\beta}$ about $h_{4-t^2, T-t^2}(Z_{t^2})$. Since $h_{4-t^2, T-t^2}(\delta i) \notin \mathcal{B}$, the result follows. $\square$

Lemma 5.6. *For all subpower functions ψ, ϕ , there is a subpower function $\bar{\psi}$ such that if T is ψ -good at δ , then the following holds for all $\delta \leq t \leq 2$. Suppose*

$$t \phi(1/t)^{-1} \leq y \leq t \phi(1/t), \quad (5.16)$$

$$|x| \leq [t + \delta] \phi(1/t). \quad (5.17)$$

Then

$$t^{-\beta} \bar{\psi}(1/t)^{-1} \leq \left| h'_{4-t^2, T-t^2}(x + iy) \right| \leq t^{-\beta} \bar{\psi}(1/t).$$

Proof. By Proposition 2.3 and (5.10) and (5.11),

$$\tilde{\psi}(1/t)^{-1} |h'_{4-t^2, T-t^2}(Z_{t^2})| \leq \left| h'_{4-t^2, T-t^2}(x + iy) \right| \leq \tilde{\psi}(1/t) |h'_{4-t^2, T-t^2}(Z_{t^2})|,$$

where $Z_{t^2} = Z_{t^2}(i\delta)$. The result then follows from (5.13). $\square$

Lemma 5.7. *For all subpower functions ψ, ϕ there is a subpower function $\bar{\psi}$ such that if T is ψ -good at δ , then the following holds for $1 \leq t \leq 2$. Suppose $w = x + iy$ with*

$$\delta \leq y \leq 1, \quad (x/y)^2 + 1 \leq \phi(1/y).$$

Then,

$$y^{-\beta} \bar{\psi}_\delta(1/y)^{-1} \leq |h'_{t^2}(w)| \leq y^{-\beta} \bar{\psi}_\delta(1/y).$$

In particular,

$$y^{-\beta} \bar{\psi}_\delta(1/y)^{-1} \leq |h'_{t^2}(iy)| \leq y^{-\beta} \bar{\psi}_\delta(1/y). \quad (5.18)$$

Proof. We will do the case $t = 2$; the argument is similar for $1 \leq t \leq 2$. We let $\bar{\psi}$ denote a subpower function in this proof, but its value may change from line to line. Since $(x/y)^2 + 1 \leq \phi(1/y)$, we can see from Proposition 2.3 that

$$\bar{\psi}(1/y)^{-1} |h'_{t^2}(iy)| \leq |h'_{t^2}(w)| \leq \bar{\psi}(1/y) |h'_{t^2}(iy)|,$$

so we may assume $x = 0$. Let $s = y$. Using the Loewner equation (5.2), we can see that there is a $c < \infty$ such that

$$y \leq \text{Im} h_{s^2}(iy) \leq cy,$$

$$|\text{Re} h_{s^2}(iy)| \leq cy,$$

$$|h'_{s^2}(iy)| \leq c.$$

The last estimate and (5.8) imply that

$$|h'_4(y)| \asymp |h_{4-s^2, T-s^2}(Z_{s^2}(iy))|.$$

Using (5.14), we see that

$$|\text{Re} Z_{s^2}(iy)| \leq y [c + \psi(1/y)].$$

By the previous result,

$$\overline{\psi}(1/y)^{-1} t^{-\beta} \leq |h'_{4-s^2, T-s^2}(Z_{s^2}(iy))| \leq \overline{\psi}(1/y) t^{-\beta}$$

□

Definition. If n is a positive integer and $j = 1, \dots, n^2$, we say that (j, n) is ψ -good if $T = 1 + (j-1)n^{-2}$ is ψ -good at n^{-1} . We let $E_{j,n}$ denote the event “ (j, n) is ψ -good” and $E(j, n)$ denotes the indicator function of $E_{j,n}$.

It is important to note that on the event $E_{j,n}$, (5.18) implies that (4.9), the corresponding estimate for $|\hat{f}'|$, holds, with perhaps a different choice of subpower function ψ . The main estimate for the lower bound is the following.

Proposition 5.8. *If $r < r_c$, here exists a subpower function ψ and $c > 0$ such that for all n and all $j = 1, 2, \dots, n^2$,*

$$\mathbb{E} \left[|\hat{f}'_{j,n}(i/n)|^\lambda E_{j,n} \right] \geq c n^{-\zeta}. \quad (5.19)$$

Remark. For fixed n , the expectation in (5.19) is the same for all j .

We will not include a proof of Proposition 5.8 because it has essentially appeared in [9, Theorem 10.8]; see also [5, Lemma 4.4]. We point out that the assumption $r < r_c$ is crucial for the result. The proof in [9] and [5] uses a careful analysis of a relatively simple one-dimensional diffusion.

5.3. Correlations. In this subsection we fix a subpower function ψ such that Proposition 5.8 holds. If n is a positive integer, we write j, k for positive integers satisfying $1 \leq j < k \leq n^2$. We will consider $E_{j,n} \cap E_{k,n}$ with indicator function $E(j, n) E(k, n)$. If j, k, n are fixed we write

$$S = 1 + \frac{j-1}{n^2}, \quad T = 1 + \frac{k-1}{n^2},$$

$$\tilde{h}_t = h_{t,S}, \quad h_t = h_{t,T},$$

and recall that this means that $\tilde{h}_t, h_t$ are solutions to the reverse-time Loewner equation with $V_{S-t} - V_S$ and $V_{T-t} - V_T$ as driving functions, respectively.

Proposition 5.9. *There is a subpower function ϕ such that for all $1 \leq j < k \leq n^2$*

$$\mathbb{E} \left[|\hat{f}'_T(i/n)|^\lambda |\hat{f}'_S(i/n)|^\lambda E(j, n) E(k, n) \right] \leq n^{-2\zeta} \left(\frac{n^2}{k-j} \right)^{\frac{\lambda\beta+\zeta}{2}} \phi \left(\frac{n^2}{k-j} \right).$$

Moreover, on the event $E_{j,n} \cap E_{k,n}$,

$$|\hat{f}_T(i/n) - \hat{f}_S(i/n)| \geq \left(\frac{k-j}{n^2} \right)^{\frac{1-\beta}{2}} \phi \left(\frac{n^2}{k-j} \right)$$

Proof. We write ϕ for a subpower function but we let its value vary from line to line; in the end we choose the maximum of all the subpower functions mentioned. Recall that $\hat{f}'_S(i/n) = \tilde{h}'_S(i/n)$, $\hat{f}'_T(i/n) = h'_T(i/n)$ and

$$\hat{f}_T(i/n) - \hat{f}_S(i/n) = \tilde{h}_S(Z_{T-S}(i/n)) - \tilde{h}_S(i/n).$$

The second assertion of the proposition follows immediately from Lemma 5.5, so we need only show the first.

Since T is ψ -good at $1/n$, we know from (5.13) that

$$|h'_T(i/n)| \leq |h'_{T-S}(i/n)| \left(\frac{n^2}{k-j} \right)^{\beta/2} \phi \left(\frac{n^2}{k-j} \right).$$

Therefore,

$$\begin{aligned} & \mathbb{E} \left[|\hat{f}'_T(i/n)|^\lambda |\hat{f}'_S(i/n)|^\lambda E(j, n) E(k, n) \right] \leq \\ & \left(\frac{n^2}{k-j} \right)^{\lambda\beta/2} \phi \left(\frac{n^2}{k-j} \right) \mathbb{E} \left[|h'_{T-S}(i/n)|^\lambda |\tilde{h}'_S(i/n)|^\lambda E(j, n) E(k, n) \right]. \end{aligned}$$

Note that $|h'_{T-S}(i/n)|^\lambda E(k, n)$ and $|\tilde{h}'_S(i/n)|^\lambda E(j, n)$ are independent random variables. Therefore,

$$\begin{aligned} & \mathbb{E} \left[|h'_{T-S}(i/n)|^\lambda |\tilde{h}'_S(i/n)|^\lambda E(j, n) E(k, n) \right] \leq \\ & \mathbb{E} \left[|\tilde{h}'_S(i/n)|^\lambda E(j, n) \right] \mathbb{E} \left[|h'_{T-S}(i/n)|^\lambda E(k, n) \right]. \end{aligned}$$

We now apply Theorem 4.1 to see that the right hand side above is bounded above by

$$n^{-\zeta} (k-j)^{-\zeta/2} \phi \left(\frac{n^2}{k-j} \right) = n^{-2\zeta} \left(\frac{n^2}{k-j} \right)^{\zeta/2} \phi \left(\frac{n^2}{k-j} \right),$$

and this concludes the proof. $\square$

6. PROOF OF THEOREM 3.3

In this section we will use the *forward* Loewner flow to prove Theorem 3.3, which we restate for the convenience of the reader.

Theorem. *If $0 < \kappa < 8$ and $1/2 \leq \alpha \leq \alpha_*$, then with probability one there exists a set V such that $\dim_h[\gamma(V)] \leq F_{\text{tip}}(\alpha)$ and for $t \notin V$, $\gamma(t) \in \mathbb{H}$,*

$$\tilde{\mu}(t, 2^{-n}) \preceq 2^{-n\alpha}, \quad n \rightarrow \infty.$$

Throughout we will fix $\kappa = 2/a < 8$. We will write u rather than α (to avoid having both α and a in formulas). To prove the theorem it suffices to show that for every bounded domain $D \subset \mathbb{H}$ bounded away from the real line, there is a set V_D with $\dim_h[\gamma(V_D)] \leq F_{\text{tip}}(u)$ and such that (3.9) holds for $t \notin V_D$ with $\gamma(t) \in D$. We fix such a D and allow constants to depend on D . The basic strategy is typical for establishing upper bounds for multifractal spectra. We estimate a particular moment of $|g'_\tau(z)|$ for an appropriate stopping time, use Chebyshev's inequality to get an estimate on

probabilities, and use this estimate to bound the dimension of a well chosen covering.

We warn the reader again that some of the notation in this section is not consistent with that in other sections.

We parametrize SLE_κ so that the conformal maps g_t satisfy

$$\partial_t g_t(z) = \frac{a}{g_t(z) - U_t}, \quad g_0(z) = z, \quad (6.1)$$

where $U_t = -B_t$ is a standard Brownian motion. This is valid for $z \in \mathbb{C} \setminus \{0\}$ up to time $T_z \in (0, \infty]$. We let H_t be the unbounded component of $\mathbb{H} \setminus \gamma(0, t]$.

6.1. Preliminaries. Let

$$Z_t = Z_t(z) = X_t + iY_t = g_t(z) - U_t.$$

If $z \in \mathbb{H}$, let

$$\Delta_t = |g'_t(z)|, \quad \Upsilon_t = \frac{Y_t}{|g'_t(z)|}, \quad \Theta_t = \arg Z_t, \quad S_t = \sin \Theta_t.$$

Υ_t equals $1/2$ times the conformal radius of H_t about z (or we can think of it as the conformal radius normalized so that the conformal radius of $\mathbb{H}$ about i equals 1). The Koebe-1/4 theorem implies that

$$\frac{\Upsilon_t}{2} \leq \text{dist}(z, \partial H_t) \leq 2\Upsilon_t. \quad (6.2)$$

Straightforward computations using (6.1) show that for $z \in \mathbb{H}$,

$$\partial_t \Delta_t = \Delta_t \frac{a(Y_t^2 - X_t^2)}{|Z_t|^4}, \quad \partial_t \Upsilon_t = -\Upsilon_t \frac{2aY_t^2}{|Z_t|^4}.$$

There exists $0 < c_1 < c_2 < \infty$ such that for $z \in D$,

$$c_1 \leq \Upsilon_0 \leq c_2, \quad c_1 \leq S_0 \leq 1. \quad (6.3)$$

Let for $u \geq 1/2$

$$r = r(u) = \frac{1}{2} - 2a - \frac{1}{4u - 2}, \quad (6.4)$$

$$\lambda = \lambda_r = \frac{r^2}{2a} + r \left(1 - \frac{1}{2a}\right), \quad \xi = \xi_r = \frac{r^2}{4a} = \frac{\lambda}{2} - \frac{r}{2} \left(1 - \frac{1}{2a}\right). \quad (6.5)$$

Note that r increases with u . Define

$$\hat{u}_c = \frac{1}{2} + \frac{1}{8a - 2}.$$

Note that $\hat{u}_c < \alpha_* = 2a/(4a - 1)$. If $u < \hat{u}_c$, since $a > 1/4$,

$$r < r(\hat{u}_c) = 1 - 4a < \min \left\{ \frac{1}{2} - 2a, 2 - 3a \right\},$$

$$r < 0, \quad r + \lambda > 0.$$

The following is a straightforward Itô's formula calculation that we omit.

Proposition 6.1. *Suppose $r \in \mathbb{R}$ and λ, ξ are as in (6.5). If $z \in \mathbb{H}$, and*

$$M_t = M_t(z) = |Z_t|^r Y_t^\xi \Delta_t^\lambda = S_t^{-r} \Upsilon_t^{\xi+r} \Delta_t^{\lambda+r},$$

then M_t is a local martingale satisfying

$$dM_t = M_t \frac{r X_t}{|Z_t|^2} dB_t.$$

Let $\mathcal{D}_n$ denote the of dyadic rationals in $\mathbb{C}$

$$z = \frac{j}{2^n} + i \frac{k}{2^n}, \quad j, k \in \mathbb{Z}.$$

Note that if $w \in \mathbb{C}$, then there exists $z \in \mathcal{D}_n$ with $|z - w| \leq 2^{-n}$ and hence $\mathcal{B}(w, 2^{-n}) \subset \mathcal{B}(z, 2^{-n+1})$.

6.2. Basic strategy. Let

$$\tau_{n,z} = \inf \{s : \Upsilon_s(z) \leq 2^{-n+3}\}.$$

We will only consider n sufficiently large so that $2^{-n+4} \leq c_1$ where c_1 is the constant in (6.3). Note that $\mathbb{P}\{\tau_{n,z} = \infty\} > 0$. If $\tau_{n,z} < \infty$, (6.2) implies

$$2^{-n+2} \leq \text{dist}(z, \partial H_{\tau_{n,z}}) = \text{dist}(z, \gamma(0, \tau_{n,z}]) \leq 2^{-n+4}.$$

In particular, if $|w - z| \leq 2^{-n}$, $\text{dist}(w, \partial H_{\tau_{n,z}}) \geq 2^{-n+1}$.

Recall that we defined the normalized harmonic measure

$$\text{hm}_t(V) = \lim_{y \rightarrow \infty} y \text{hm}(iy, V, H_t)$$

and $\tilde{\mu}(t, \epsilon) = \text{hm}_t(\overline{\mathcal{B}}(\gamma(t), \epsilon))$. Similarly we define

$$\hat{\text{hm}}_t(V) := \lim_{y \rightarrow \infty} y \text{hm}(iy, \partial V, H_t \setminus V),$$

and note that $\text{hm}_t(V) \leq \hat{\text{hm}}_t(V)$. Set $\text{hm}_{n,z} = \text{hm}_{\tau_{n,z}}$ and similarly for $\hat{\text{hm}}$. If $|z - \gamma(t)| \leq 2^{-n}$, then $\tau_{n,z} \leq t$ and hence by monotonicity of harmonic measure,

$$\hat{\text{hm}}_{n,z}[\mathcal{B}(z, 2^{-n+1})] \geq \text{hm}_t[\mathcal{B}(z, 2^{-n+1})] \geq \tilde{\mu}(t, 2^{-n}).$$

Let $\mathcal{D}_n(D)$ denote the set of $z \in \mathcal{D}$ such that $\text{dist}(z, D) \leq 2^{-n}$ and

$$A_m^u = A_m^u(D) = \bigcup_{n=m}^{\infty} \bigcup_z \mathcal{B}(z, 2^{-n+1}),$$

where the inner union is over all $z \in \mathcal{D}_n(D)$ satisfying

$$\hat{\text{hm}}_{n,z}[\mathcal{B}(z, 2^{-n+1})] \geq 2^{-nu}. \quad (6.6)$$

Then if $\gamma(t) \in D \setminus A_m^u$, for all n sufficiently large

$$\tilde{\mu}(t, 2^{-n}) \leq 2^{-nu}.$$

Hence for each m , A_m^u is a cover of $D \cap V_u$ where V_u is the set of $\gamma(t)$ that do not satisfy (3.9). Let $N_n = N_{n,u}(D)$ be the cardinality of the set of $z \in \mathcal{D}_n(D)$ satisfying (6.6). Then for all s ,

$$\mathcal{H}^s[D \cap V_u] \leq \lim_{m \rightarrow \infty} \mathcal{H}^s(A_m^u) \leq c \lim_{m \rightarrow \infty} \sum_{n=m}^{\infty} N_n 2^{-ns}.$$

The following proposition follows immediately.

Proposition 6.2. *Suppose $u, s > 0$ and*

$$N_{n,u}(D) \preceq 2^{ns}, \quad n \rightarrow \infty.$$

Then

$$\dim_h [D \cap V_u] \leq s.$$

Proof. The argument above shows that for all $s' > s$, $\mathcal{H}^{s'}[D \cap V_u] = 0$. $\square$

In order to show that with probability one for SLE that

$$\dim_h [D \cap V_u] \leq s.$$

it suffices to show that

$$\mathbb{E}[N_{n,u}(D)] \preceq 2^{ns}, \quad n \rightarrow \infty.$$

Indeed, this relation and the Borel-Cantelli lemma imply that with probability one for all $s' > s$, $N_{n,u}(D) \leq 2^{ns'}$. Note that

$$\mathbb{E}[N_{n,u}(D)] \leq c_D 2^{2n} \sup_{\text{dist}(z,D) \leq 2^{-n}} \mathbb{P} \left\{ \tau_{n,z} < \infty; \hat{\text{hm}}_{n,z}[\mathcal{B}(z, 2^{-n+1})] \geq 2^{-nu} \right\}.$$

Notice that conformal invariance of harmonic measure and distortion estimates imply that on the event $\tau_{n,z} < \infty$,

$$\hat{\text{hm}}_{n,z}[\mathcal{B}(z, 2^{-n+1})] \asymp 2^{-n} |g'_{\tau_{n,z}}(z)|.$$

Indeed, $g_{\tau_{n,z}}(\mathcal{B}(z, 2^{-n+1}))$ is a connected set whose diameter is comparable to $2^{-n} |g'_{\tau_{n,z}}(z)|$ and whose distance from the real axis is comparable to $2^{-n} |g'_{\tau_{n,z}}(z)|$. Hence, there exists $c < \infty$ such that

$$\mathbb{E}[N_{n,u}(D)] \leq c_D 2^{2n} \sup_{\text{dist}(z,D) \leq 2^{-n}} \mathbb{P} \left\{ \tau_{n,z} < \infty; |g'_{\tau_{n,z}}(z)| \geq c 2^{-n(u-1)} \right\}.$$

In the remainder of this section we will show that there exists $c = c_D < \infty$ such that for all n sufficiently large and all z with $\text{dist}(z, D) \leq 2^{-n}$,

$$\mathbb{P} \left\{ \tau_{n,z} < \infty; |g'_{\tau_{n,z}}(z)| \geq c 2^{-n(u-1)} \right\} \leq 2^{-n\rho(u)}, \quad (6.7)$$

where

$$\rho(u) = \left\lceil \frac{1}{8a} + 2a - 1 \right\rceil \left(u - \frac{1}{2}\right) + \left\lceil \frac{1}{2} - \frac{1}{8a} \right\rceil + \frac{1}{32a(u - \frac{1}{2})}.$$

Then from the above arguments we know that with probability one,

$$\dim_h [D \cap V_u] \leq 2 - \rho(u) = F_{\text{tip}}(u).$$

The second equality is a straight-forward calculation. The remainder of this section is devoted to establishing (6.7).

6.3. Weighting by the martingale. The local martingale M_t is not a martingale because it “blows up” on the event of measure zero that z is on the path $\gamma(0, \infty)$. However, if we choose stopping times τ such as $\tau_{n,z}$ which prevent the path from getting too close to z , then the stopped process $M_{t \wedge \tau}$ is a martingale. Let $\mathbb{P}^*, \mathbb{E}^*$ denote probabilities and expectations with respect to the measure obtained by weighting by (the stopped martingale) M . The Girsanov theorem implies that

$$dB_t = \frac{r X_t}{|Z_t|^2} dt + dW_t, \quad 0 \leq t < \tau,$$

where W_t is a standard Brownian motion with respect to the measure $\mathbb{P}^*$. In particular,

$$d\Theta_t = \frac{(1 - 2a - r) X_t Y_t}{|Z_t|^4} dt - \frac{Y_t}{|Z_t|^2} dW_t.$$

It is useful to use a “radial” parametrization $\sigma(t)$. We write $\hat{Z}_t = Z_{\sigma(t)}$, $\hat{X}_t = X_{\sigma(t)}$, etc. The radial parametrization is defined by

$$\hat{\Upsilon}_t := \Upsilon_{\sigma(t)} = e^{-2at}.$$

Note that

$$-2a \hat{\Upsilon}_t = \partial_t \hat{\Upsilon}_t = -2a \frac{\hat{Y}_t^2}{|\hat{Z}_t|^4} \partial_t \sigma(t),$$

which implies

$$\partial_t \sigma(t) = \frac{|\hat{Z}_t|^4}{\hat{Y}_t^2}.$$

Note also that

$$d\hat{\Theta}_t = (1 - 2a) \cot \hat{\Theta}_t dt + d\hat{B}_t,$$

and the local martingale M_t satisfies

$$d\hat{M}_t = -r \hat{M}_t \cot \hat{\Theta}_t d\hat{B}_t.$$

Moreover, we have that

$$d\hat{\Theta}_t = (1 - 2a - r) \cot \hat{\Theta}_t dt + d\hat{W}_t. \quad (6.8)$$

In the above, $\hat{B}_t$ and $\hat{W}_t$ are standard Brownian motions with respect to $\mathbb{P}$ and $\mathbb{P}^*$ respectively. Since $1 - 2a - r > 1/2$, we compare with a Bessel process to see that in the measure $\mathbb{P}^*$, $\hat{\Theta}_t$ never reaches $\{0, \pi\}$, see [7, Chapter 1]. It follows that $\hat{M}_t$ is actually a martingale. Also, the invariant probability density for the SDE (6.8) equals

$$f(\theta) = c \sin^{2(1-2a-r)} \theta.$$

Since $r < 1 - 4a < 3 - 4a$ it follows that $\sin^r$ is integrable with respect to $f(\theta) d\theta$. The important fact for us, is that there is a c such that if $\hat{\Theta}_t$ satisfies (6.8) with $\sin \hat{\Theta}_0 \geq c_1$, then for all $t > 0$,

$$\mathbb{E}^*[\hat{S}_t^r] \leq c. \quad (6.9)$$

Let

$$\tau_s = \inf \{t : \Upsilon_t = e^{-2as}\}.$$

For $r < \frac{1}{2} - 2a$, we have for all s ,

$$\mathbb{P}^*\{\tau_s < \infty\} = 1.$$

Then, using (6.9),

$$\begin{aligned} \mathbb{E} \left[|g'_{\tau_s}(z)|^{\lambda+r}; \tau_s < \infty \right] &= \mathbb{E} \left[M_{\tau_s} S_{\tau_s}^r \Upsilon_{\tau_s}^{-\xi-r}; \tau_s < \infty \right] \\ &= e^{2as(\xi+r)} \mathbb{E} \left[M_{\tau_s} S_{\tau_s}^r; \tau_s < \infty \right] \\ &= e^{2as(\xi+r)} M_0(s) \mathbb{E}^* \left[\hat{S}_s^r \right] \\ &\leq c e^{2as(\xi+r)}. \end{aligned}$$

Since $\lambda + r > 0$, if $\epsilon = e^{-2as}$,

$$\mathbb{P} \left\{ \tau_s < \infty; |g'_{\tau_s}(z)| \geq \epsilon^{u-1} \right\} \leq \epsilon^{-(u-1)(\lambda+r)} \mathbb{E}[|g'_{\tau_s}(z)|^{\lambda+r}; \tau_s < \infty] \leq c \epsilon^{\rho(u)}$$

where

$$\rho(u) = -(u-1)(\lambda+r) - (r+\xi).$$

Doing the algebra, we get

$$\rho(u) = \left[\frac{1}{8a} + 2a - 1 \right] \left(u - \frac{1}{2} \right) + \left[\frac{1}{2} - \frac{1}{8a} \right] + \frac{1}{32a(u - \frac{1}{2})}.$$

This proves (6.7) which concludes the proof of Theorem 3.3. □

REFERENCES

- [1] V. Beffara (2008). The dimension of the SLE curves, *Annals of Probab.* **36**, 1421–1452.
- [2] D. Beliaev, S. Smirnov (2009). Harmonic measure and SLE, to appear in *Comm. Math. Phys.*
- [3] B. Duplantier (2004). Conformal fractal geometry & boundary quantum gravity in *Fractal Geometry and Applications: A jubilee of Benoît Mandelbrot*, M. Lapdus & M. van Frankenhuijsen, ed., AMS, 365–482.
- [4] I. Binder, B. Duplantier (2002), Harmonic measure and winding of conformally invariant curves. *Phys. Rev. Lett.* **89**, 264101.
- [5] F. Johansson Viklund, G. Lawler (2010), Optimal Hölder exponent for the SLE path, to appear in *Duke Math. J.*
- [6] N.-G. Kang (2007). Boundary behavior of SLE, *Journal of AMS*, **20**, 185–210.
- [7] G. Lawler (2005). *Conformally Invariant Processes in the Plane*, Amer. Math. Soc.
- [8] G. Lawler (2009). Schramm-Loewner evolution, in *Statistical Mechanics*, S. Sheffield and T. Spencer, ed., IAS/Park City Mathematical Series, AMS, 231–295.

- [9] G. Lawler (2009). Multifractal analysis of the reverse flow for SLE, in *Fractal Geometry and Stochastics IV*, C. Brandt, P. Mörters, & M. Zähle, ed., Birkhäuser, 73–107.
- [10] G. Lawler and S. Sheffield (2010), The natural parametrization for the Schramm-Loewner evolution, to appear in *Ann. Probab.*
- [11] G. Lawler, O. Schramm, W. Werner (2001). Values of Brownian intersection exponents I: half-plane exponents, *Acta Math.* **187**, 237–273.
- [12] G. Lawler, O. Schramm, W. Werner (2001). Values of Brownian intersection exponents II: plane exponents, *Acta Math.* **187**, 275–308.
- [13] G. Lawler, O. Schramm, W. Werner (2002). Values of Brownian intersection exponents III: two-sided exponents, *Ann. Inst. Henri Poincaré.* **38**, 109–123.
- [14] G. Lawler, O. Schramm, W. Werner (2004). Conformal invariance of planar loop-erased random walks and uniform spanning trees, *Ann. Probab.* **32**, 939–995.
- [15] J. Lind (2008). Hölder regularity of the *SLE* trace, *Trans. AMS* **360**, 3557–3578.
- [16] N. G. Makarov (1998), Fine structure of harmonic measure. *Algebra i Analiz* 10 (1998), no. 2, 1–62; translation in *St. Petersburg Math. J.* 10 (1999), no. 2, 217–268.
- [17] Ch. Pommerenke (1992), *Boundary behaviour of conformal maps*. Springer-Verlag, Berlin, 1992.
- [18] S. Rohde and O. Schramm (2005). Basic properties of SLE, *Annals of Math.* **161**, 879–920.
- [19] O. Schramm (2000). Scaling limits of loop-erased random walks and uniform spanning trees, *Israel J. Math.* **118**, 221–288.
- [20] S. Smirnov (2001). Critical percolation in the plane: conformal invariance, Cardy’s formula, scaling limits, *C. R. Acad. Sci. Paris Sr. I Math.*, **333**, 239–244.
- [21] S. Smirnov (2010) Conformal invariance in random cluster models. I. Holomorphic fermions in the Ising model, to appear in *Ann. Math.* **172**.
- [22] W. Werner (2004), Random planar curves and Schramm-Loewner evolutions, *Ecole d’Eté de Probabilités de Saint-Flour XXXII - 2002*, *Lecture Notes in Mathematics* **1840**, Springer-Verlag, 113–195.

FREDRIK JOHANSSON VIKLUND, DEPARTMENT OF MATHEMATICS, COLUMBIA
UNIVERSITY, 2990 BROADWAY, NEW YORK, NY 10027, USA
E-mail address: `fjv@math.columbia.edu`

GREGORY F. LAWLER, DEPARTMENT OF MATHEMATICS AND DEPARTMENT OF
STATISTICS, UNIVERSITY OF CHICAGO, 5734 S. UNIVERSITY AVENUE, CHICAGO, IL
60637, USA
E-mail address: `lawler@math.uchicago.edu`